

SYSTEMATIC MAP PROTOCOL

Open Access



Outcomes of wildlife translocations in protected areas: what is the type and extent of existing evidence? A systematic map protocol

Joseph Langridge^{*} , Romain Sordello and Yorick Reyjol

Abstract

Background: Conversion, fragmentation, and loss of natural habitats are among the main causes of declining species' populations worldwide. Protected areas are therefore crucial for biodiversity as they provide refuge and ensure key ecological processes. Wildlife translocations, defined as "the deliberate movement of organisms from one site for release in another", have been used in conjunction as a conservation tool for a number of decades as wild populations become increasingly fragmented and endangered. Not only are translocations used to bolster the viability of imperiled species but are also recommended for improving population resilience and adapting species' ranges in response to climate change. Despite translocation being a recognised conservation tool, it remains complex with variable results due to the different factors that can determine its success. Accordingly, the Map will investigate the existing evidence on the links between different types of wildlife translocation interventions and factors that may be important to consider for planning. This will provide an overview of relevant studies for possible future syntheses, and may help to inform management decisions.

Method: We will perform a thorough search of peer-reviewed journal articles and grey literature sources documenting the occurrence of translocations in the context of protected areas. Two databases will be used: Web of science core collection and Scopus, with a supplementary search in Google Scholar. Multiple key specialized websites will also be used. All bibliographic data will be extracted, managed, and screened in Microsoft excel. Three screening stages will be undertaken (title, then abstract, then full texts) against predefined inclusion criteria. The retained relevant literature will be subjected to coding and meta-data extraction. No formal validity appraisal will be undertaken. The Map will particularly highlight translocation operations in terms of origin and destination (i.e. translocating from one protected area to another, within the same area, and from and to non-protected areas) by taxonomic group, among other important factors (e.g. number of individuals, age class, release strategy, distance between capture and release sites etc.). Finally, a database will be provided along with a Map narratively describing the evidence with summary figures and tables of pertinent study characteristics.

Keywords: Managed relocations, Reintroduction, Supplementation, Introduction, Conservation areas

Background

Modification, fragmentation, and loss of natural habitats are among the main causes of declining species' populations worldwide [1–4]. In the face of such threats, extinction rates have been accelerating and biological diversity diminishing for the last several decades [5–7]. Accordingly, protected areas such as national parks, nature and

*Correspondence: joseph.langridge1@mnhn.fr
UMS Patrimoine Naturel (PatriNat), OFB-CNRS-MNHN, 36 Rue
Geoffroy-Saint-Hilaire CP41, 75005 Paris, France



© The Author(s) 2020. This article is licensed under a Creative Commons Attribution 4.0 International License, which permits use, sharing, adaptation, distribution and reproduction in any medium or format, as long as you give appropriate credit to the original author(s) and the source, provide a link to the Creative Commons licence, and indicate if changes were made. The images or other third party material in this article are included in the article's Creative Commons licence, unless indicated otherwise in a credit line to the material. If material is not included in the article's Creative Commons licence and your intended use is not permitted by statutory regulation or exceeds the permitted use, you will need to obtain permission directly from the copyright holder. To view a copy of this licence, visit <http://creativecommons.org/licenses/by/4.0/>. The Creative Commons Public Domain Dedication waiver (<http://creativecommons.org/publicdomain/zero/1.0/>) applies to the data made available in this article, unless otherwise stated in a credit line to the data.

biosphere reserves play vital roles in maintaining refuge and ensuring fundamental ecological mechanisms such as dispersal and gene exchange [5, 8–11]. Further, not only do they provide key habitat and conserve biodiversity from various human pressures [8, 10] but they are able to maintain higher species population levels, including threatened species, better than other management approaches [10, 11].

Translocation is an umbrella term referring to the “deliberate movement of organisms from one site for release in another” [12]. Indeed, it may occur in different contexts such as reintroduction in which organisms are transported and released into their historical native range but from which they have become extirpated or extinct, or supplementation (also known as reinforcement), which refers to the addition of individuals to an existing population of conspecifics [12]. Thirdly, introduction, which from a conservation perspective is often referred to as assisted colonization [12–14], attempts to establish a species outside of its recorded historical distribution but within appropriate habitat and biogeographical area. Each strategy sharing the ultimate goal of population persistence [14].

Historically, the intentional movement and release of species has occurred for millennia [12, 15], but the use of translocations to address species-focused conservation objectives is more recent [15]. For example, between 1973 and 1989 more than 700 translocations were estimated to have taken place per year across the USA, Australia, and New Zealand in order to restore and enhance populations [5]. Latterly, interest in assisted colonization has been driven by predicted habitat and climate changes [13, 15]. In 1985, Peters and Darling [16] suggested that climate change might alter habitat suitability for species confined within protected areas, effectively stranding them as habitat becomes increasingly unfavourable [15]. They proposed the translocation of individuals into new reserves encompassing habitat that was or would become appropriate [16], therefore potentially compensating for the fixed nature of perimeters [17, 18]. More recently, the concept of rewilding has emerged [15, 19]. Originally based on the keystone role played by wide-ranging predators and their ability to maintain ecosystem equilibrium through top-down trophic interactions [15], the concept has since harmonized with the current conservation translocation framework to include the role of species reintroduction to restore ecological processes [15, 19], and to a broader extent, the restoration of ecosystem functions by means of introducing ecological replacements [15].

In the current context of the biodiversity crisis, translocations and particularly reintroductions of threatened species are more numerous [20]. They are also used in

conjunction with conservation areas more regularly as populations become progressively more fragmented and endangered [21]. Even though past efforts have not been entirely uniform with a notably marked taxonomic bias towards birds and mammals (e.g. [5, 22]) and an apparent prioritization for larger more charismatic species [15, 22, 23], attention being paid to other groups has rapidly increased since the early 2000s (e.g. English Nature’s Species Recovery Programme involving 62 species, of which only 11 were birds or mammals) [22]. Thus, for management purposes, the need to synthesize this profuse information is apparent. Moreover, concerning success, studies regularly identify the value of habitat quality at recipient sites and the importance of species being relocated to non-degraded habitats [5, 20, 24, 25]. Indeed, in a previous review on plant reintroductions, Godefroid et al. [26] confirmed that reintroducing species to protected areas significantly increased survival rate. Equally, regarding vertebrate translocations, several papers highlight the positive effects of protected habitat (e.g. [27]), and in a number of canid translocations protected areas were regularly chosen as release sites [28–30]. With the overarching pressure of climate change, several authors have continued to propose translocations as a viable means to enhance the resilience of threatened species, improve ecosystem integrity, and assist migration to favourable habitats [9, 14, 24, 31, 32].

Despite the number of translocations rapidly growing and it increasingly being recognized as a key conservation measure, implementation is often complex and different programs have had varying results. From a biological perspective, this is notably due to the numerous different factors that influence its success [33, 34] such as the number of translocated individuals [35], the distances involved [36], whether acclimatisation strategies (e.g. protective enclosures or supplemental feeding) are used [37], and what levels of habitat quality individuals are faced with at release sites [25, 26]. From a social perspective, interventions are still considered controversial: cost, feasibility, and political acceptability remain the principal influencing factors [38].

Although previous overviews exist (e.g. [23, 34]), and while others have explored the effectiveness of anti-predator training and conditioning interventions (e.g. [37]), there appears to be a deficit in terms of systematic literature assessments on the role of protected areas. Hence, our aim is to map evidence of translocation operations carried out in the context of protected areas detailing the distribution and abundance of relevant studies in relation to key factors that influence success. This will provide an evidence base for possible future reviews, and should

help to inform eventual management and stakeholder decisions.

Stakeholder engagement

The current systematic map will be conducted as part of a wider European LIFE programme (the EU's funding instrument for environment and climate action). The LIFE project, entitled "Natur'Adapt", is coordinated by the French Nature Reserves Network (*Les Réserves Naturelles de France* (RNF)), and co-financed by the French Ministry of Ecology and the French Office of Biodiversity (OFB). RNF is accompanied by nine other beneficiaries, including The French National Natural History Museum (*Muséum National d'Histoire Naturelle* (MNHN)), who will be responsible for the mapping process.

The project's principal aim is to align conservation efforts in protected areas to the challenges associated with climate change, in France and across Europe. Subsequently, the progressive development of an adaptation plan will be undertaken based, firstly, on six "experimental" nature reserves then progressively made adaptable to all protected areas in France and Europe. The MNHN is responsible for a key LIFE action: to provide evidence syntheses. This will help reserve managers build their adaptation plan by transferring scientific knowledge to them in an accessible and summarized form. As a first step, several working groups were conducted between RNF, MNHN, and reserve managers. This was an opportunity for reserve managers to define all relevant conservation strategies, in the context of climate change, of which they were most in need of scientific evidence to support decision-making. At the end of this process, translocation, among other measures, was retained as it was considered a necessary conservation action plan. As a result of numerous discussions, a systematic map was chosen as a central reference tool. Further workshops were held to specifically learn the stakeholders' needs and involve them in the definition of the Map's meta-data variables.

Objectives of the review

The main objective is to systematically map translocation operations within the context of protected areas (i.e. operations from, to or within a protected area). The IUCN protected area management categories will be used for this as they represent a global standard for defining conservation areas. In agreement with the specific aims of the LIFE project, this Map will consider translocations for species conservation—where the primary goal is to improve the status of the focal species through supplementation, reintroduction, or assisted migration. Translocation for rewilding—where the initial

motive is to restore natural ecosystem functions will be included. In accordance with Seddon et al. [15], translocation rewilding will only entail (i) population restoration through reintroduction, where release within the indigenous range aims at reestablishing some ecological function, or (ii) in the form of a conservation introduction through ecological replacement [15, 19]. Neither invasive species nor historical introductions for hunting purposes will be included. We will aim to provide a comprehensive overview of the distribution of studies by taxonomic group and type of translocation, in conjunction with other key drivers (e.g. age class, release strategy, distance between capture and release sites, number of individuals initially translocated etc.) that may influence various biological outcomes i.e. success of wildlife translocation operations.

Therefore, the primary question for this Map protocol is as follows: What type, extent, and distribution of evidence exists on the outcomes of wildlife translocations carried out in protected areas?

Components of the primary question in Table 1.

Methods

Searching for articles

Our search strategy has been designed in order to retrieve a broad range of articles covering the topic of wildlife translocations in protected areas. Indeed, the systematic map will follow the Environmental Evidence Guidelines and conforms to the ROSES standards (see Additional file 1 for our declaration and checklist of adherence to the ROSES guidelines).

Search terms and languages

All searches will be performed using English terms only. Hence, all relevant studies published in English will be included in this systematic map. This will include diverse bibliographic documents (e.g. books conference proceedings, journal articles, theses, technical reports etc.)

Search strings

Firstly, a scoping exercise was conducted in the Web of Science Core Collection database to explore the efficiency of chosen words and the number of articles returned. In accordance with our main objective, we combined all search terms relating to protected areas and wildlife translocations. Concerning protected areas, the chosen key words represent synonyms of the different types of reserves and management categories that exist.

Thus, the search string that produced the highest efficiency is presented below (see Additional file 2 for test list details and Additional file 3 for information of the building process of the search string).

Table 1 Components of the systematic map

Population (P)	All plant and animal species of wild or captive source			
Intervention (I)	<i>Type of interventions</i>	<i>Definition of interventions</i>	<i>Conservation aim of intervention</i>	
	Introduction	This refers to the intentional manual transfer/movement and release of an organism outside of its indigenous range/historical distribution [12]	<p>(i) <i>Assisted migration</i>: this refers to the intentional manual transfer/movement and release outside of the indigenous range, to primarily avoid extinction of populations of the focal species [12]</p> <p>(ii) <i>Ecological replacement</i>: This refers to the intentional manual transfer/movement and release of an organism outside its indigenous range/historical distribution, to perform a specific ecological function [12]</p>	This occurs if the persistence of a species in its indigenous range is threatened from current or future impacts than at alternative sites [12]
	Reintroduction	This refers to the intentional manual transfer/movement and release of an organism inside its indigenous range/historical distribution but from which it has disappeared or become extinct locally, regionally, or otherwise (No conspecifics are present in situ) [12]		The conservation aim is to re-establish a viable population of the focal species within its historical range [12]
	Supplementation	This refers to the intentional manual transfer/movement and release of an organism into the existing distribution of a population of conspecifics [12]		The aim is to enhance and reinforce population viability e.g. by increasing population size, or by increasing genetic diversity [12]
Comparator (C)	Studies will not be required <i>stricto sensu</i> to have a comparator. Although in certain cases the study design may translate as a time series comparison (before and after translocation)			
Outcomes (O)	All relevant outcomes related to the translocated population, including space use, demography, survival, reproduction, feeding, behaviour, genetics, and physiology (<i>cf.</i> Table 3 for full typology)			
Context (C) ^a	<i>Type of protected areas</i>	<i>Definitions of protected areas</i>		
	Strict reserves for the protection of nature (Ia)	Areas set aside to strictly protect biodiversity where human visitation, use, and impacts are strongly limited [39]		
	Wilderness areas (Ib)	Areas that are largely unmodified, retaining their natural character, and free of inappropriate or excessive human use or presence [39]		
	National Parks (II)	Protected areas of large natural or near natural areas set aside to protect large-scale ecological processes [39].		
	Natural monuments (III)	Protected areas set aside to protect a specific natural feature in the landscape [39].		
	Management areas (IV)	Specific protected areas that aim to safeguard a particular species or habitat. Consequently, the management reflects this priority [39]		
	Protected landscapes (V)	A protected area where humans and nature together over time have produced an area of significant ecological, biological, cultural and scenic value [39]		
Protected areas with sustainable use of natural resources (VI)	Protected areas which conserve ecosystems and habitats together with associated cultural values and traditional natural resource management and use [39]			

^a Concerning context, this will equate to all interventions from, to, or within these types of protected areas

TS=(“protected area\$” OR “protected landscape\$” OR “protected site\$” OR “receptor site\$” OR “reintroduction site\$” OR “natur* reserve\$” OR “national park\$” OR “regional park\$” OR “national reserve\$” OR “biological reserve\$” OR “biosphere reserve\$” OR “regional reserve\$” OR “wilderness area\$” OR “natural monument\$” OR “management area\$” OR sanctuar*) AND TS=(“assisted colonization” OR “assisted population migration” OR “assisted migration” OR “assisted gene flow” OR “managed relocation\$” OR transloc* OR reintroduc* OR reinforc* OR “assisted range expansion\$” OR “assisted long-distance migration\$” OR rewilding OR “wild release”).

Estimating the comprehensiveness of the search

A test list of 40 scientific articles was established and used to assess the comprehensiveness of the search string. The test list was composed of relevant scientific articles identified by the review team prior to the mapping process. The overall comprehensiveness was 100%. [Two additional files provide further details (see Additional files 2 and 3)].

Publication databases to be searched

All published material will be collected from the following databases (and managed in excel).

- Web of Science (WOS) core collection. The entire database i.e. all citation indexes will be searched by Topic i.e. using the “TS” field tag, which searches for key words in the title, abstract and key-words of published documents (see Additional file 4 for Web of Science subscription details).
- Scopus. We will equally search for all published documents. We will use the field tag “TITLE-ABS-KEY”, which operates in the same way as the “TS” tag in WOS.

These databases were chosen as they provide comprehensive citation data for numerous different academic disciplines. The English search string detailed above will be used for both literature sources. The search string will be adapted as necessary to account for the differences in the use of field tags and Boolean characters [an additional file provides details on number of search hits and dates of searches (see Additional file 5)].

Internet searches to be conducted

A supplementary retrieval of publications will be undertaken using web-based search engines.

- Google Scholar (<https://scholar.google.com/>). We used the same key words in the software programme Publish or perish (version 6) to retrieve all academic citations. The software’s use of Boolean characters differs from WOS and Scopus. As a result, the search string was broken down into eight separate searches, in order to achieve a similar comprehensiveness, as only a single term can be included in the field “all of the words”. Consequently, each sub-search was limited to the first 200 search hits, in line with recommendations [40]. (Refer to Additional file 5.)
- A retrieval of theses will also be done using UK Theses and Dissertations (<https://ethos.bl.uk>). We will search for theses using the intervention key words only. We will search using five key words: “reintroduction” OR “reinforcement” OR “introduction” OR “translocation” OR “rewilding”. Hits limited to 200.
- Conservation Evidence (<https://www.conservationalevidence.com/>)—we will collect primary research using the Journal’s “Advanced search”. Use of five key words: “reintroduction”, “reinforcement”, “introduction”, “translocation”, or “rewilding” will be used for collecting individual studies. We will extract the first 40 hits per keyword search (total hits: 200).

Specialist searches

The following specialist organisations will be searched for reports which contain translocations to, from and within protected areas.

- US Federal Science database (<https://www.science.gov/>).
- US Fish and wildlife service (<https://www.fws.gov/>).
- Office National de la Chasse et de la Faune Sauvage (<http://www.oncfs.gouv.fr/>).
- IUCN Conservation Planning Specialist Group’s document library (<http://www.cpsg.org/document-repository>).
- Association of Zoos & Aquariums (<https://www.aza.org/>). Including the European Association (<https://www.eaza.net/>).
- Rewilding Europe (<https://rewilding-europe.com/>).

Supplementary searches

A call for literature will be made through the professional networks of *Les Réserves Naturelles de France* (RNF) and EuroParc. An advert will be

published in the monthly newsletter of RNF. Euro-Parc, who act as a federation of protected areas at the European continental scale will also solicit their network. Since translocations programs are often carried out without being published in the form of scientific articles this will provide further opportunity to gather additional grey literature such as PhD and MSc theses, various technical reports, and other documentation. AirTable, which works like a database will be the specific software used to acquire the documents sent via the stakeholders' contacts.

Article screening and study eligibility criteria

Screening process

In accordance with the pre-defined screening and study eligibility criteria (detailed in “[Eligibility criteria](#)” section), study selection will follow a three-stage filtering process carried out by two members of the mapping team. Firstly, all titles will be screened, followed by abstracts and thirdly full texts. During screening, we will choose to take a conservative approach. Hence, if the qualifying information is not detailed sufficiently to reject or to retain with certainty, then the article in question will be kept for assessment at the next eligibility stage in the overall filtering process. In addition, articles or grey literature that qualify after title screening but do not contain an abstract will pass by default to the full-text screening stage. Lastly, should our search string retrieve, in addition, any relevant published material in French it will also be incorporated into the mapping process because these are the two languages spoken and understood by all members of the map team.

Consistency checking

To fully assess whether both reviewers adhere to the eligibility criteria, a Kappa test will be performed at the start of each filtering stage. Accordingly, 10% of retained titles, 10% of retained abstracts, and 10% of retained full texts will be pre-screened to check for agreement. Kappa scores should be equal to or greater than 0.6. If differences of opinion occur, the process will be repeated with new samples until a score of 0.6 or greater is reached. Even if statistical agreement is reached, all (if any) remaining disagreements will be discussed before beginning the screening process. A consistency check for meta-data extraction will also be undertaken based on training articles representing 10% of the retained corpus. All eventual disagreements will be discussed between the reviewers.

Eligibility criteria

Different eligibility criteria will be applied at each filtering stage. Table 2 describes a summary description of the eligibility criterion.

Title

Inclusion criteria Firstly, all titles will be retained if presence of the terms reintroduction, supplementation (and its common synonyms i.e. reinforcement, augmentation, re-stocking, enhancement) and introduction (and its common synonyms i.e. assisted migration, managed relocation etc.). Secondly, any title containing compatible synonyms such as, re-wilding, release, range-shifts, transfer, restoration etc., will also be retained. In cases where none of the above words are present, a title would still meet eligibility if it strongly implies a translocation event (i.e. reference to captive of wild stock) or meta-population management. *Nb.* At title screening stage, all types of literature (including review, meta-analyses and relevant discussion and opinion articles) will be retained.

Exclusion criteria clear absence of the above key words. Translocation in a genetic context, e.g. chromosomal translocation, will also be excluded.

Abstract

Inclusion criteria Presence of words related to survival, mortality, space use, genetics and all other relevant biological outcomes (*cf.* Table 3). The abstract will also be retained if it contains words confirming a translocation event to, from, within or away from protected area perimeters. Additionally, for the purpose of the Map, if the translocation event has occurred to solve human-wildlife conflicts then this will also satisfy the inclusion criteria.

Exclusion criteria If no obvious description of intervention exists.

Full text

Inclusion criteria Primarily but not exclusively, if the outcome has been obtained from field studies (e.g. individuals equipped with radio-collars at time of release, reported number of individuals surviving after a pre-determined time-scale). However, discussion and review articles will be retained if presence of PICO elements is sufficiently described. Also, if the article presents evidence of comparison of release strategies. All articles that clearly state that population/individuals (plants or animals) are of captive or wild stock and have been transferred to, from, or within protected areas.

Exclusion criteria Similar to those applied for title or abstract screening, or information informing that the translocated population is invasive or introduced

Table 2 Systematic map inclusion and exclusion criteria, and PICO definitions for the three-stage screening process

Criteria	Description	PICO definition(s)
Inclusion criteria		
Eligible population	Any terrestrial, marine, or aquatic plant or animal species of wild or captive source	IN: Wild—free-ranging species from natural environments; not domesticated or cultivated Captive—species born, bred, or cultivated in captive setting (e.g. zoos, nurseries)
Eligible intervention	Conservation-based translocation operations (i.e. reintroduction, supplementation and introduction) (see Table 1 for definitions)	IN: Conservation-based interventions—where the aim is improving viability and persistence of translocated population. Regarding introduction, this will also include ecological replacement and assisted migration [14]
Eligible comparator	Studies will not be required <i>stricto sensu</i> to have a comparator present	NA
Eligible outcomes	All relevant biological outcomes related to the subject population, including space use, demography, survival, reproduction, trophic, behaviour, genetics, and physiology	IN: For full description and definitions (<i>cf.</i> Table 3)
Exclusion criteria		
Ineligible population	Invasive species or population historically introduced for hunting purposes	OUT: Invasive alien species outside of its natural distribution area and that threatens biological function and diversity of native populations Game/Hunting purposes: if transfer of species is for hunting purposes
Ineligible intervention	Non conservation-based introductions (e.g. historically introduced animals for hunting purposes or accidental introductions)	OUT: Non-conservation based transfer of species—where persistence or conservation of individuals is not the motivation (e.g. sport/game hunting) Accidental introductions: any documentation reporting on accidental invasions, pathways, and risks etc.
Ineligible outcomes	Sources not studying one of either Space use, Demography, Survival, Reproduction, Feeding, Behaviour, Genetics, Physiology	OUT: Non-relevant outcomes that would not enable any interpretation of success. Trophic cascades will not be retained

Table 3 Outcome categories and corresponding descriptions

Outcome category	Description	Example references (from test list or scoping exercise)
Space use	Studies measuring all movement/dispersal of translocated individuals. This will include notably home range measurements, or euclidean distance travelled	[43]
Demography	Studies outlining the changes in number of individuals, males/females, of the translocated population i.e. population growth overtime	[44, 45]
Survival	Studies illustrating precisely the proportion of individuals alive or level of mortality since translocation	[46]
Reproduction	Any impacts on reproduction, expressed by number of young born since translocation, or specifically the survival rate of offspring	[47]
Feeding	All impacts specifically on diet and feeding of translocated individuals. (<i>Nb.</i> cascade effects will not be included as an outcome)	[48, 49]
Behaviour	Studies measuring changes in terms of communication (e.g. vocal), social structure, or anti-predator behaviour i.e. stress/vigilance levels, of translocated individuals	[50, 51]
Genetics	Studies relating to the genetic structure of the translocated species	[52]
Physiology	All biological or physiological impacts measured at the molecular, cellular or organic level (e.g. hormone activity)	[53]

for hunting purposes. We will provide a list of articles excluded at full text with reasons for exclusion.

Study validity assessment

No formal validity appraisal of included studies will be performed. All studies that are deemed eligible at the full text stage based on the Population, Intervention, Outcomes, and Context criteria/screening stages will be included in the Map. Thus, this Systematic Map will be considered a thorough narrative synthesis ahead of any review providing a comprehensive and robust overview of the existing evidence.

Data coding strategy

A thorough meta-data extraction for the Map will be performed by the same two members of the mapping team. Each selected article will be double coded. If, due to resource limitations, true double coding is not possible, a posteriori cross-check will be carried out and potential disagreements will be discussed until a consensus is reached. Concerning missing data, if data is not sufficiently detailed or simply unknown, it will be coded as such. The following meta-data will be extracted from all articles retained after completion of the screening process [an additional file is provided with full explanations (see Additional file 6)]:

Bibliographic information

- Authors of article.
- Title and abstract.
- Year of publication.

- Publication source (name of journal).
- Full-text language (English, French or other).
- Document type (journal article, book, conference object, thesis (Phd, or Msc), technical documentation, or other).
- Study content (study, review, meta-analysis, discussion paper, modelling, or other).

Study characteristics

- Study country.
- Capture and release site locality coded as two separate fields (name and geographic coordinates will be recorded if given).
- Capture and release site climate types coded as two separate fields (under the Köppen-Geiger Climate Classification).
- IUCN protected area management categories coded for each protected area. (This will be achieved by accessing the IUCN PAs database via <http://www.protectedplanet.net>, and then matching each PA with the PAs in the database based on NAME).
- Protected area (this will be coded in order to decipher if individuals are translocated from-to different protected areas, to, from, or within same protected area).
 - From-to: transfer from one protected area (PA) to another.
 - To: transfer from a non-protected habitat i.e. outside of PA perimeter to a PA.

- iii. From: transfer from a PA to a non-protected habitat i.e. outside of PA perimeter.
 - iv. Within same: transfer occurring within the same PA perimeter.
- Study area biome (as stated by article authors). But regrouped into 6 broad categories (Additional file 6 gives explanations on chosen habitat classes):
 - i. Forest/wooded.
 - ii. Savannah.
 - iii. Open habitats.
 - iv. Wetland/humid.
 - v. Marine.
 - vi. Aquatic.
 - Study release strategy. Two release strategies will be coded as follows:
 - i. Soft release: studies having sufficiently described methods to acclimatize individuals at the recipient site. Two key methods will define soft release: use of protective enclosures, and use of supplemental feeding [41, 42].
 - ii. Hard release: immediate release (no acclimatization and no supplementary food) [41].
 - Study cost (in the rare case that such information is reported, we will record figures stated by article authors).
 - Distance between capture and release site (coded *a posteriori* with recorded geographic coordinates and use of geographic software).

Population characteristics

- The lowest taxonomic rank will be recorded if sufficiently detailed i.e. species name. Otherwise, higher taxonomic classification will be used e.g. genus, family or Order.
- Source and destination (wild-to-wild, captive-to-wild, breeding-to-wild). If transferred individuals are bred at specific sites, then released to wild this will be coded as “breeding-to-wild”.
- Study sample size (number of individuals initially translocated, as stated by article authors).
- Study age class at release (adult, juveniles or both, as stated by authors). However, concerning plant translocations it will be appropriate to detail life stage at translocation e.g. seed, seedling, and adult plant.

Intervention characteristics

- Study interventions (supplementation, reintroduction, or introduction). 5 possible intervention categories will be coded as follows:
 - i. Introduction—if a study is based on a single one-off intervention i.e. assisting the migration of a given species to suitable habitat outside of its historical distribution.
 - ii. Intro + Suppl—an introduction intervention followed for the supplementation of the same introduced population.
 - iii. Reintroduction—a single one-off reintroduction event (not followed by supplementation).
 - iv. Reintro + Suppl—a reintroduction followed by the supplementation of the same reintroduced population.
 - v. Supplementation—where a given study only reports on the supplementation of an already threatened species.
- Duration of intervention i.e. “translocation period” (number of years). This will be relevant for cases where an initial reintroduction event is followed by several supplementations.
- If translocation is climate motivated or not.
- Programme motivation (this will outline the overall motive of the manual transfer/movement of the species in question).
 - i. Conservation (improving status of focal species).
 - ii. Rewilding (restoring natural functions).
 - iii. Experimental or trial translocations.
 - iv. Human-wildlife conflict.
 - v. Wildlife rescue operations (from human development projects/urbanisation).
 - vi. Metapopulation management.

Outcome characteristics

- The following biological outcomes will be recorded: space use, demography, survival, reproduction, feeding, behaviour, genetics, and physiology (*cf.* Table 3 for full descriptions).

Study mapping and presentation

A systematic map database will be provided, detailing all included articles from the full text screening stage. The

systematic map will include all the metadata coded for each article. For the cases where more than one study is reported in the same article, each study will be recorded as a unique entry in the excel database with its corresponding geographical coordinates, if given, and a unique study ID. This database will be available as an open access excel spreadsheet and included as an appendix to the systematic map publication.

The map database will be described in the map publication with summary figures and tables of the relevant study characteristics. A geographic map will present the location of each translocation event/study. Possible knowledge gaps (under-represented subtopics that warrant further primary research) and knowledge clusters (well-represented subtopics for full synthesis by a systematic review) will be identified by cross-tabulating key meta-data variables (e.g. biological groups and outcomes). Based on these results, recommendations will be made on priorities for future research concerning translocation and protected areas. Recommendations will also be made to inform management. To this end, regarding the specific objectives of the LIFE project, all Map results will be transferred to reserve managers. In addition, a practitioner brief will be provided to reserve managers with the aim of summarizing key results in an operational manner in order to aid decision making. Workshops are already planned for this.

Supplementary information

Supplementary information accompanies this paper at <https://doi.org/10.1186/s13750-020-00199-4>.

Additional file 1. Our declaration and checklist of adherence to the ROSES guidelines.

Additional file 2. Test list and overall comprehensiveness of search string.

Additional file 3. Illustrating the building process of the search string.

Additional file 4. Web of Science Core Collection database subscription details.

Additional file 5. Corresponding number of Search hits from Web of Science core collection, Scopus, and supplementary search in google scholar and organisation websites.

Additional file 6. A Codebook outlining the meta-data extraction methods.

Acknowledgements

We would like to thank the reserve managers of *Les Réserves Naturelles de France* for their contribution during the round-tables. We thank François Sarrazin of the Sorbonne University and Bruno Colas of the University of Paris-Sud for their expertise on the subject. We would also like to thank Dakis-Yaoba Ouédraogo for her invaluable feedback.

Authors' contributions

JL and RS wrote the protocol. JL, RS and YR conceived the literature search strategy including the key word equation. RS contributed to the initial round-table and integration of the systematic review methods into the LIFE project, and provided overall assistance concerning methods and CEE guidelines. YR

provided overall scientific expertise and proof-reading. All authors read and approved the final manuscript.

Funding

This work will be undertaken within the framework of the LIFE program entitled "Natur'Adapt". The project is co-financed by the LIFE programme (European commission), The French Ministry of Ecology and the French Office of Biodiversity.

Availability of data and materials

Data sharing is not applicable to the publication of the review protocol. All datasets associated with the Systematic Map will be made available as open access files (Additional file 6).

Ethics approval and consent to participate

No ethics agreement was required.

Consent for publication

No consent for publication was required.

Competing interests

Authors declare having no competing interests.

Received: 29 January 2020 Accepted: 17 July 2020

Published online: 29 July 2020

References

- Murphy SE, Greenaway F, Hill DA. Patterns of habitat use by female brown long-eared bats presage negative impacts of woodland conservation management. *J Zool*. 2012;288:177–83.
- Ceballos G, Ehrlich PR, Barnosky AD, Garcia A, Pringle RM, Palmer TM. Accelerated modern human-induced species losses: entering the sixth mass extinction. *Sci Adv*. 2015;1:e1400253.
- Nowakowski AJ, Thompson ME, Donnelly MA, Todd BD. Amphibian sensitivity to habitat modification is associated with population trends and species traits. *Glob Ecol Biogeogr*. 2017;26:700–12.
- Newbold T, Hudson LN, Hill SLL, Contu S, Lysenko I, Senior RA, et al. Global effects of land use on local terrestrial biodiversity. *Nature*. 2015;520:45–50.
- Griffith B, Scott J, Carpenter J, Reed C. Translocation as a species conservation tool: status and strategy. *Science*. 1989;245:477–80.
- Caizergues A, Rätti O, Helle P, Rotelli L, Ellison L, Rasplu J-Y. Population genetic structure of male black grouse (*Tetrao tetrix* L.) in fragmented vs. continuous landscapes. *Mol Ecol*. 2003;12:2297–305.
- Rivera-Ortiz FA, Aguilar R, Arizmendi MDC, Quesada M, Oyama K. Habitat fragmentation and genetic variability of tetrapod populations. *Anim Conserv*. 2015;18:249–58.
- Cantú-Salazar L, Gaston KJ. Very large protected areas and their contribution to terrestrial biological conservation. *Bioscience*. 2010;60:808–18.
- Lunt ID, Byrne M, Hellmann JJ, Mitchell NJ, Garnett ST, Hayward MW, et al. Using assisted colonisation to conserve biodiversity and restore ecosystem function under climate change. *Biol Conserv*. 2013;157:172–7.
- Watson JEM, Dudley N, Segan DB, Hockings M. The performance and potential of protected areas. *Nature*. 2014;515:67–73.
- Gray CL, Hill SLL, Newbold T, Hudson LN, Börger L, Contu S, et al. Local biodiversity is higher inside than outside terrestrial protected areas worldwide. *Nat Commun*. 2016;7:1–7.
- IUCN. Guidelines for reintroductions and other conservation translocations. IUCN. 2013. <https://www.iucn.org/content/guidelines-reintroductions-and-other-conservation-translocations>. Accessed 31 Oct 2013.
- Hoegh-Guldberg O, Hughes L, McIntyre S, Lindenmayer DB, Parmesan C, Possingham HP, et al. Assisted colonization and rapid climate change. *Science*. 2008;321:345–6.
- Seddon PJ. From reintroduction to assisted colonization: moving along the conservation translocation spectrum. *Restor Ecol*. 2010;18:796–802.
- Seddon PJ, Griffiths CJ, Soorae PS, Armstrong DP. Reversing defaunation: restoring species in a changing world. *Science*. 2014;345:406–12.

16. Peters RL, Darling JDS. The Greenhouse Effect and Nature Reserves. Global warming would diminish biological diversity by causing extinctions among reserve species. *BioScience*. 1985;35:707–17.
17. Mawdsley JR, O'Malley R, Ojima DS. A review of climate-change adaptation strategies for wildlife management and biodiversity conservation. *Conserv Biol*. 2009;23:1080–9.
18. McLachlan JS, Hellmann JJ, Schwartz MW. A framework for debate of assisted migration in an era of climate change. *Conserv Biol*. 2007;21:297–302.
19. Sandom CJ, Dempsey B, Bullock D, Ely A, Jepson P, Jimenez-Wisler S, et al. Rewilding in the English uplands: policy and practice. *J Appl Ecol*. 2019;56:266–73.
20. Sarrazin F, Barbault R. Reintroduction: challenges and lessons for basic ecology. *Trends Ecol Evol*. 1996;11:474–8.
21. Evans K, Moore R, Harris S. The social and ecological integration of captive-raised adolescent male African elephants (*Loxodonta africana*) into a wild population. *PLoS ONE*. 2013;8:e55933.
22. Hodder KH, Bullock JM. Translocations of native species in the UK: implications for biodiversity. *J Appl Ecol*. 1997;34:547–65.
23. Hale SL, Koprowski JL. Ecosystem-level effects of keystone species reintroduction: a literature review. *Restor Ecol*. 2018;26:439–45.
24. Seddon PJ, Armstrong DP, Maloney RF. Developing the science of reintroduction biology. *Conserv Biol*. 2007;21:303–12.
25. McCoy ED, Osman N, Hauch B, Emerick A, Mushinsky HR. Increasing the chance of successful translocation of a threatened lizard. *Anim Conserv*. 2014;17:56–64.
26. Godefroid S, Piazza C, Rossi G, Buord S, Stevens A-D, Aguraiuja R, et al. How successful are plant species reintroductions? *Biol Cons*. 2011;144:672–82.
27. Müller J, Wölfel M, Wölfel S, Müller DWH, Hothorn T, Heurich M. Protected areas shape the spatial distribution of a European lynx population more than 20 years after reintroduction. *Biol Cons*. 2014;177:210–7.
28. Moehrenschrager A, Somers M. Canid reintroductions and metapopulation management. Cambridge: IUCN/SSC Canid Specialist Group, Gland; 2004. p. 289–97.
29. Vogel JT, Somers MJ, Venter JA. The foraging ecology of reintroduced African wild dog in small protected areas. *wbio*. Nordic Board for Wildlife Research; 2018. <https://bioone.org/journals/Wildlife-Biology/volume-2018/issue-1/wlb.00424/The-foraging-ecology-of-reintroduced-African-wild-dog-in-small-10.2981/wlb.00424.full>. Accessed 2 Aug 2020.
30. Ripple WJ, Beschta RL. Wolf reintroduction, predation risk, and cottonwood recovery in Yellowstone National Park. *For Ecol Manage*. 2003;184:299–313.
31. Heller NE, Zavaleta ES. Biodiversity management in the face of climate change: a review of 22 years of recommendations. *Biol Cons*. 2009;142:14–32.
32. Prober SM, Doerr VAJ, Broadhurst LM, Williams KJ, Dickson F. Shifting the conservation paradigm: a synthesis of options for renovating nature under climate change. *Ecol Monogr*. 2019;89:e01333.
33. Goldenberg SZ, Owen MA, Brown JL, Wittemyer G, Oo ZM, Leimgruber P. Increasing conservation translocation success by building social functionality in released populations. *Global Ecol Cons*. 2019;18:e00604.
34. Fischer J, Lindenmayer DB. An assessment of the published results of animal relocations. *Biol Cons*. 2000;96:1–11.
35. Brambell MR. Reintroduction. *Int Zool Yearbook*. 1977;17:112–6.
36. Brown JR, Bishop CA, Brooks RJ. Effectiveness of short-distance translocation and its effects on western rattlesnakes. *J Wildlife Manag*. 2009;73:419–25.
37. Tetzlaff SJ, Sperry JH, DeGregorio BA. Effects of antipredator training, environmental enrichment, and soft release on wildlife translocations: a review and meta-analysis. *Biol Cons*. 2019;236:324–31.
38. Corlett RT. Restoration, reintroduction, and rewilding in a changing world. *Trends Ecol Evol*. 2016;31:453–62.
39. Dudley N. Guidelines for applying protected area management categories. IUCN; 2008. <https://portals.iucn.org/library/node/9243>. Accessed 15 Jun 2020.
40. Haddaway NR, Collins AM, Coughlin D, Kirk S. The role of Google Scholar in evidence reviews and its applicability to grey literature searching. *PLoS ONE*. 2015;10:e0138237.
41. de Milliano J, Stefano JD, Courtney PR, Temple-Smith PD, Coulson G. Soft-release versus hard-release for reintroduction of an endangered species: an experimental comparison using eastern barred bandicoots (*Perameles gunnii*). *Wildl Res*. 2016;43:12.
42. Broughton S, Dickman C. The effect of supplementary food on home range of the southern brown bandicoot, *Isodon obesulus* (Marsupialia: Peramelidae). *Aust J Ecol*. 2006;16:71–8.
43. Fernando P, Leimgruber P, Prasad T, Pastorini J. Problem-elephant translocation: translocating the problem and the elephant? *PLoS ONE*. 2012;7:e50917.
44. Hochkirch A, Agnes W, Anje T, Friedhelm N. Translocation of an endangered insect species, the field cricket (*Gryllus campestris* Linnaeus, 1758) in northern Germany. *Biodivers Conserv*. 2007;16:3597–607.
45. Willis SG, Hill JK, Thomas CD, Roy DB, Fox R, Blakeley DS, et al. Assisted colonization in a changing climate: a test-study using two UK butterflies. *Cons Lett*. 2009;2:46–52.
46. Islam MZ, Ismail K, Boug A. Restoration of the endangered Arabian Oryx *Oryx leucoryx*, Pallas 1766 in Saudi Arabia lessons learnt from the twenty years of re-introduction in arid fenced and unfenced protected areas. *Zool Middle East*. 2011;54:125–40.
47. Bodinof CM, Briggler JT, Junge RE, Mong T, Beringer J, Wanner MD, et al. Survival and body condition of captive-reared juvenile ozark hellbenders (*Cryptobranchus alleganiensis bishopi*) following translocation to the wild cope. *Am Soc Ichthyol Herpetol*. 2012;2012:150–9.
48. Clayton J, Pavey C, Vernes K, Jefferys E. Diet of mala (*Lagorchestes hirsutus*) at Ulu-ru-Kata Tju-ta National Park and comparison with that of historic free-ranging mala in the Tanami Desert: implications for management and future reintroductions. *Aust Mammal*. 2015;37:201–11.
49. Stannard HJ, Caton W, Old JM. The diet of red-tailed phascogales in a trial translocation at Alice Springs Desert Park, Northern Territory, Australia. *J Zool*. 2010;280:326–31.
50. Renan S, Speyer E, Ben-Nun T, Ziv A, Greenbaum G, Templeton AR, et al. Fission-fusion social structure of a reintroduced ungulate: implications for conservation. *Biol Cons*. 2018;222:261–7.
51. Viljoen JJ, Ganswindt A, Reynecke C, Stoeger AS Jr. Vocal stress associated with a translocation of a family herd of African elephants (*Loxodonta africana*) in the Kruger National Park, South Africa. *Bioacoustics*. 2015;24:1–12.
52. Hicks JF, Rachlow JL, Rhodes OE, Williams CL, Waits LP. Reintroduction and genetic structure: rocky Mountain Elk in Yellowstone and the Western States. *J Mammal*. 2007;88:129–38.
53. Schultz D, Rich B, Rohrig W, McCarthy P, Mathews B, Corrigan A, et al. Investigations into the health of brush-tailed rock wallabies (*Petrogale penicillata*) before and after reintroduction. *Aust Mammal*. 2011;33:152–61.

Publisher's Note

Springer Nature remains neutral with regard to jurisdictional claims in published maps and institutional affiliations.

SYSTEMATIC MAP

Open Access



Existing evidence on the outcomes of wildlife translocations in protected areas: a systematic map

Joseph Langridge, Romain Sordello*  and Yorick Reyjol

Abstract

Background: Ecosystem degradation, mainly through overexploitation and destruction of natural habitats, is a well-known threat to the viability and persistence of many species' populations worldwide. The use of translocations as a viable conservation tool in conjunction with protected areas has been rapidly increasing over the last few decades. Protected areas such as strict nature reserves, national parks, and species management areas continue to be central tools for biodiversity conservation as they provide vital habitats set aside from various human pressures. Because action consistently runs ahead of policy, the need for a clearer evidence base on the outcomes of wildlife translocations undertaken at a global scale is becoming increasingly urgent for scientific and decision-making communities, in order to build clear strategy frameworks around conservation translocations. We therefore conducted a systematic mapping exercise to provide an overview of the existing evidence on the outcomes of wildlife translocations in protected areas.

Methods: We searched two bibliographic databases, four web-based search engines with search-by-key-words capacity, 5 specialist websites, and conducted a grey literature call through two project stakeholders. We screened articles by title, abstract, and full text using pre-defined inclusion criteria all the while assessing the consistency of the reviewers. All relevant translocations were coded from retained publications. Key variables of interest were extracted and coded for each translocation event. The quantity and characteristics of the available evidence and knowledge gaps/clusters are summarised. The distribution and frequency of translocations are presented in heat- and geographical maps.

Review findings: A total of 613 articles were considered eligible for coding bibliometric data. Metapopulation management and review articles were not coded for quantitative and qualitative variables. Linked data (duplicated translocations) were also excluded. Finally, 841 studies of different translocation events were fully coded from 498 articles. Most of these translocations were carried out in North America and Oceania. The most commonly undertaken intervention types were one-off supplementations and "supplemented reintroductions". Mammals were by far the most transferred group among animals. *Magnoliopsida* was the most translocated plant group. Survival, space use, and demography metrics were the most studied outcomes on translocated species.

Conclusions: This systematic map provides an up-to-date global catalogue of the available evidence on wildlife translocations to, from, or within protected areas. It should enable protected area managers to better understand their role in the global network of protected areas, regarding translocation practice, both as suppliers or recipients of

*Correspondence: romain.sordello@mnhn.fr
UMS PatriNat OFB-MNHN-CNRS, 36 rue Geoffroy-Saint-Hilaire CP41,
75005 Paris, France



© The Author(s) 2021. **Open Access** This article is licensed under a Creative Commons Attribution 4.0 International License, which permits use, sharing, adaptation, distribution and reproduction in any medium or format, as long as you give appropriate credit to the original author(s) and the source, provide a link to the Creative Commons licence, and indicate if changes were made. The images or other third party material in this article are included in the article's Creative Commons licence, unless indicated otherwise in a credit line to the material. If material is not included in the article's Creative Commons licence and your intended use is not permitted by statutory regulation or exceeds the permitted use, you will need to obtain permission directly from the copyright holder. To view a copy of this licence, visit <http://creativecommons.org/licenses/by/4.0/>. The Creative Commons Public Domain Dedication waiver (<http://creativecommons.org/publicdomain/zero/1.0/>) applies to the data made available in this article, unless otherwise stated in a credit line to the data.

translocated species. It may help managers and practitioners make their own choices by comparing previous experiences, regarding both the species concerned and the precise translocation modalities (number of individuals, etc.). Finally, it constitutes a decision-making tool for managers as well as for policy makers for future translocations.

Keywords: Managed relocations, Reintroduction, Reinforcement, Supplementation, Ecological replacements, Assisted migration, Rewilding, Conservation areas

Background

Ecosystem degradation, mainly through overexploitation and destruction of natural habitats, is a well-known threat to the viability and persistence of many species' populations worldwide [1–5]. According to the IUCN's Global Species Programme, more than 35,000 species across the globe are currently threatened with extinction [6]. Today, species extinction rates are 100 to 1000 times greater than the natural background level, with populations of wild species declining at a rate of 0.5 to 1% per annum on average [7–11]. Indeed, the major causes of this global biodiversity crisis are anthropogenic in nature, and exacerbated by ongoing climate changes [10]. However, protected areas (i.e. "a clearly defined geographical space, recognised, dedicated and managed, through legal or other effective means, to achieve the long-term conservation of nature [...] [12]) such as wilderness areas, national parks, and habitat or species management areas continue to be central tools for biodiversity conservation as they provide vital habitats, set aside from various human pressures for the safeguard of species [13–15]. Furthermore, especially in fragmented environments, the role of protected areas is paramount as they provide higher quality habitat often maintaining higher species population levels than other management strategies [13, 16, 17].

Historically, the intentional movement and release of species has occurred for millennia [18, 19], but the use of translocations to address species-focused conservation objectives is a more recent intervention. With several authors advocating the importance of release site habitat quality [e.g. 20–23], the implementation of "intentional movements of organisms from one site for release in another" [18] in conjunction with protected areas may be an increasingly essential conservation strategy. With the overriding pressure of climate change expected to alter habitat suitability for species confined within current protected areas—effectively stranding them as habitat becomes increasingly unfavourable [24]—several authors suggest intentionally moving species between protected areas to where appropriate habitats are currently, or predicted to persist [24–26].

Indeed, different forms of conservation translocations exist, all of which are increasingly recognised as viable means to enhance the resilience of threatened species, improve ecosystem integrity, and assist migration to favourable habitats [25, 27–30]. Firstly, "reintroductions": aim to re-establish viable populations of a focal species within their historical range but from which they have become extirpated or extinct [15]. Secondly, "supplementations": aim to enhance and reinforce population viability by increasing population size and genetic diversity of existing resident populations [18]. Thirdly, "assisted migration": occurs if the persistence of a species in its indigenous range is threatened from current or future impacts than at alternative sites outside of its historical range [15]. In addition, the concept of rewilding has been discussed in the context of climate change [19, 31, 32]. The concept was originally based on the keystone role of apex predators and their ability to maintain ecosystem equilibrium through top-down trophic interactions [19]. However, it now includes the role of species reintroductions in order to specifically restore ecological processes or to reestablish an ecological function lost through extinction by means of introducing a suitable related taxa capable of performing a similar ecological role [19]. Further, by reintroducing ecosystem engineers, it has been suggested that this in turn can enhance the albedo effect and contribute to a greater carbon capture and overall negative feedback to global warming [32].

As the above mentioned conservation translocations are being increasingly employed in both terrestrial and marine environments [22, 33, 34], especially reintroductions of threatened species [35–38], the need to summarise this profuse information is ever more apparent for management purposes. In addition, because national and regional protected areas are frequent sites of population restoration action, given that long-term protection from diverse anthropogenic pressures have maintained relatively high-quality habitat [39–42], reserve and wildlife managers alike may be increasingly confronted with the challenges associated with such interventions, i.e. as donor or recipient site managers.

Despite the growing number of these interventions, implementation remains complex and different

programs have had varying outcomes, which is notably due to the different factors that influence success. From a biological perspective, this may include the level of habitat quality (or protection) at recipient sites [7, 22, 33, 43, 44], the number of translocated individuals [45–48], the distances involved [49], whether acclimatisation measures are used etc. [50, 51]. From a social perspective, cost, feasibility, and political acceptability are commonly cited [52]. Translocation can also be used to reduce human-wildlife conflicts because some interaction between humans and fauna and the resultant retaliatory killings can endanger the populations of these animals (felids, bears, snakes, etc.) [53, 54]. For all these reasons, it is essential to assess effectiveness of translocations in order to aid decision making. However, even before being able to perform concrete quantitative reviews (e.g. meta-analyses) on the efficacy of wildlife translocations, there is an equal need to extensively catalogue all types of wildlife translocation operations in relation to such factors providing a preliminary evidence base or a ‘landscape of knowledge’, hence the scope of this map.

Although previous overviews exist [45, 51, 55, 56], to our knowledge there are a distinct lack of literature reviews of the systematic-map nature aiming to catalogue the number of different interventions implemented to, from, and within protected areas. We analysed publications dealing with all kinds of animal and plant species translocations, worldwide. The key aims of the systematic map were to identify (1) where all types of translocations take place; (2) what the purposes and motivations are behind translocations; (3) what taxonomic groups are involved across all types of translocations; (4) what the most common direction of movement is in terms of protected areas (and what level of protection they have); (5) what distances are involved for the associated taxonomic groups; (6) how many individuals are translocated; and (7), what types of outcomes are assessed. Lastly, the financial costs of the translocations were summarised if these data were available.

Stakeholder engagement

Given the mounting evidence on the negative effects of climate change on species viability and ecosystem integrity, there is an emerging consensus within the scientific and policy communities that conservation translocation may be a necessary means to boost declining species’ populations, restore ecological processes [25], and assist species migration to favourable habitats [57]. However, more research on translocations and its risks are needed [58]. Making decisions on such a complex subject—as reserve managers may have to do—is an

arduous task because there remain uncertainties due to many different programs experiencing variable results [59]. The current work was integrated into the project “Natur’Adapt”, a European LIFE programme coordinated by the French Nature Reserves Network (RNF). Among the first actions of this LIFE, several evidence syntheses have been planned, to inform future decision-making by reserve managers. These reviews will help the reserve managers build their adaptation plan, transferring scientific knowledge in an accessible and summarized form. The selection of the evidence synthesis subjects was made in cooperation with RNF, The Natural History Museum (MNHN), and the reserve managers through dialogue and meetings. In particular, reserve managers were asked to propose the most relevant conservation/management measures in the context of climate change, of which they were most expecting information. Then, during the opening meeting of the LIFE program, a workshop was carried out with the reserve managers to select, among all propositions put forward, those that would be definitively chosen for evidence synthesis. Translocation was considered a necessary future conservation action plan in order to help align conservation efforts in protected areas to the challenges associated with climate change, in France and across Europe. A systematic map was subsequently chosen as a central reference tool that reserves managers could use as a systematised log of interventions undertaken to date.

Indeed, before being able to perform concrete quantitative syntheses on the effectiveness of wildlife translocations, RNF requested a comprehensive catalogue of existing literature describing the links between different types of wildlife translocation interventions and the factors that may be important to consider before implementation. Several further meetings were held, (refer to protocol [60] for full details), providing reserve managers the opportunity to contribute to the coding and data extraction decisions/procedures (notably the definition of variables). Following the acceptance of the protocol, further dialogue with managers enabled decisions to be made on the form and content of a practitioner brief, which will provide a summary of key results in an operational manner in order to aid decision-making.

Objective of the review

The proposed systematic map is intended to provide a thorough overview of the existing literature cataloguing the outcomes of wildlife translocations carried out in the context of protected areas (i.e. translocations where the individuals have been transferred between different protected areas, come from non-protected sites, or relocated within the same reserve). Our

primary research question was: what is the type, extent, and distribution of existing evidence on the outcomes of such wildlife translocations? The aim of this systematic map was to describe the different types of translocation conservation interventions and several factors that could be important to consider for planning (e.g., IUCN protection category, intervention type, outcomes measured, distances between capture and release sites, and taxonomic groups studied). Also, identifying knowledge gaps and knowledge clusters, which are sufficiently covered by existing studies to allow for full systematic reviewing. The systematic map was based on the following study components (see Table 1).

Methods

This systematic map followed the detailed methods described in the a priori systematic map protocol [60]. It was performed in strict concordance with the guidelines provided by The Collaboration for Environmental Evidence [61], and conforms to the ROSES reporting standards (see Additional file 1).

Deviations from the protocol

In the systematic map protocol, we had committed to performing Kappa tests on 10% of the eligible articles to assess the consistency of the reviewers during the screening process. We were able to respect these samples for titles and abstracts. Unfortunately, due to a lack of time and means, we were only able to perform our full-text Kappa test on 5% of retained articles.

Regarding the coding step (extraction of the meta-data), the systematic map protocol provided, as a first option, an independent coding for each article by two reviewers. But, in case of time constraints, it provided as a second option that an a posteriori cross-check could be carried out with discussion between the coders on potential disagreements. Indeed, due to the lack of project resources, we had to choose this alternative option.

Search for articles

Languages

Only English terms were included in the searches and, from the returned articles, those in either English and French were assessed and read, due to limited resources and the languages understood by the map team.

Search strings

The following search string was built during a scoping exercise in Web of Science, described in the systematic map protocol [60], for use in bibliographic databases (see section below):

TS=((“protected area\$” OR “protected landscape\$” OR “protected site\$” OR “receptor site\$” OR “reintroduction site\$” OR “natur* reserve\$” OR “national park\$” OR “regional park\$” OR “national reserve\$” OR “biological reserve\$” OR “biosphere reserve\$” OR “regional reserve\$” OR “wilderness area\$” OR “natural monument\$” OR “management area\$” OR “sanctuar*”) AND TS=(“assisted colonization” OR “assisted population migration” OR “assisted migration” OR “assisted gene flow” OR “managed relocation\$” OR transloc* OR reintroduc* OR reforc* OR “assisted range expansion\$” OR “assisted long-distance migration\$” OR rewilding OR “wild release”). Search strings for all the databases and searches engines are listed in Additional file 2.

Bibliographic databases

Two online bibliographic databases were searched using the search string described above:

- ‘Web of science Core Collection’ (WOSCC) accessible on the Web of Science platform (Clarivate), using the right access from the MNHN (French National Museum of Natural History). The search covered the entire database (i.e. all the citations indexes: SCI-EXPANDED, SSCI, A&HCI, CPCI-S, CPCI-SSH, BKCI-S, BKCI-SSH, ESCI and CCREXPANDED). It returned 2322 citations.
- Scopus (Elsevier), using the access from the CNRS (National Centre of Scientific Research). It returned 2563 citations.

For both the databases, the requests were run on 5 August 2020, targeting the topic (TS for WOSCC; TITLE-ABS-KEY for Scopus) and without any restriction (timespan, search area, etc.).

Web-based search engines

A supplementary retrieval of articles was undertaken using these web-based search engines:

- Google Scholar (<https://scholar.google.com/>). We used the software programme “Publish or perish (version 6)”. Given the conditions of this software (characters limit allowed), we could not use the search string described above and a selection of keywords had to be made (see Additional file 2—search strings). The software’s use of Boolean characters also differs; as a result, the search string was broken down into 10 separate sub-searches in order to achieve a similar comprehensiveness to WOSCC

Table 1 Components of the systematic map

Population (P)	All plant and animal species of wild or captive source		
Intervention (I)	<i>Type of interventions</i>	<i>Definition of interventions</i>	<i>Conservation aim of intervention</i>
	Introduction	This refers to the intentional manual transfer/movement and release of an organism outside of its indigenous range/historical distribution [13]	(i) Assisted migration: this refers to the intentional manual transfer/movement and release outside of the indigenous range, to primarily avoid extinction of populations of the focal species [13] (ii) Ecological replacement: this refers to the intentional manual transfer/movement and release of an organism outside its indigenous range/historical distribution, to perform a specific ecological function [13]
	Reintroduction	This refers to the intentional manual transfer/movement and release of an organism inside its indigenous range/historical distribution but from which it has disappeared or become extinct locally, regionally, or otherwise. (No conspecifics are present in situ) [13]	This occurs if the persistence of a species in its indigenous range is threatened from current or future impacts than at alternative sites [13] This is used to re-establish an ecological function lost through extinction; involving the most suitable existing sub-species, or a close relative of the extinct species [13]
	Supplementation	This refers to the intentional manual transfer/movement and release of an organism into the existing distribution of a population of conspecifics [13]	The conservation aim is to re-establish a viable population of the focal species within its historical range [13] The aim is to enhance and reinforce population viability e.g. by increasing population size, or by increasing genetic diversity [13]
Comparator (C)	No comparator will be required <i>stricto sensu</i> . Although in certain cases the study design may translate as a time series comparison (before and after translocation)		
Outcomes (O)	<i>Outcome category</i>	<i>Outcome description</i>	
	Space use	Studies measuring all movement/dispersal of translocated individuals. This will include notably home range measurements, or <i>Euclidean</i> distance travelled	
	Demography	Studies outlining the changes in number of individuals, males/females, of the translocated population i.e. population growth overtime	
	Survival	Studies illustrating precisely the proportion of individuals alive or level of mortality since translocation	
	Reproduction	Any impacts on reproduction, expressed by number of young born since translocation, or specifically the survival rate of offspring	
	Feeding	All impacts specifically on diet and feeding of translocated individuals. (<i>Nb.</i> cascade effects will not be included as an outcome)	
	Behaviour	Studies measuring changes in terms of communication (e.g. vocal), social structure, or anti-predator behaviour i.e. stress/vigilance levels, of translocated individuals	
	Behaviour	Studies measuring changes in terms of communication (e.g. vocal), social structure, or anti-predator behaviour i.e. stress/vigilance levels, of translocated individuals	
	Physiology	All biological or physiological impacts measured at the molecular, cellular or organic level (e.g. hormone activity)	
Context (C)^a	<i>Type of protected areas</i>	<i>Definitions of protected areas</i>	
	Strict reserves for the protection of nature (Ia)	Areas set aside to strictly protect biodiversity where human visitation, use, and impacts are strongly limited [31]	
	Wilderness areas (Ib)	Areas that are largely unmodified, retaining their natural character, and free of inappropriate or excessive human use or presence [31]	
	National Parks (II)	Protected areas of large natural or near natural areas set aside to protect large-scale ecological processes [31]	
	Natural monuments (III)	Protected areas set aside to protect a specific natural feature in the landscape [31]	
	Management areas (IV)	Specific protected areas that aim to safeguard a particular species or habitat. Consequently, the management reflects this priority [31]	
	Protected landscapes (V)	A protected area where humans and nature together over time have produced an area of significant ecological, biological, cultural and scenic value [31]	
	Protected areas with sustainable use of natural resources (VI)	Protected areas which conserve ecosystems and habitats together with associated cultural values and traditional natural resource management and use [31]	

^a Concerning context, this will equate to all interventions from, to, or within these types of protected areas

and Scopus. Each sub-search was limited to the first 200 search hits, in line with recommendations [62].

- UK Theses and Dissertations (<https://ethos.bl.uk>). We used the “Advanced search” mode and five intervention key words: reintroduction OR reinforcement OR introduction OR translocation OR rewilding. We archived the first 200 titles in an excel spreadsheet.
- Conservation Evidence (<https://www.conservationevidence.com/>). We searched for primary research papers within Journal’s “Advanced search” mode targeting individual studies only with the use of the same five key words: reintroduction, reinforcement, introduction, translocation, rewilding. Only one key word can be used at a time; we thus extracted the first 40 hits per keyword search (200 hits in total).
- US Federal Science database (<https://www.science.gov/>). We searched all literature using the topic tab “Text”, which returns reports, conference papers, and other textual information. A basic string can be used with Boolean characters. Likewise for the retrieval of theses, the string used for this organizational website was: reintroduction OR reinforcement OR introduction OR translocation OR rewilding. The First 200 hits were archived and managed in excel.

Searches on all the web-based search engines were performed on 7 July 2020 (see Additional file 2 for details about search on web-based search engines).

Specialist websites

On 27 July 2020, we conducted hand searching for relevant articles on the following six specialist websites:

- US Fish and wildlife service (<https://www.fws.gov/>).
- Office National de la Chasse et de la Faune Sauvage (<http://www.oncfs.gouv.fr/>).
- IUCN Conservation Planning Specialist Group’s document library (<http://www.cpsg.org/document-repository>).
- Association of Zoos & Aquariums (<https://www.aza.org/>). Including the European Association (<https://www.eaza.net/>).
- Rewilding Europe (<https://rewildingeuropa.com/>).

These websites do not allow a standardised literature search. Thus, the bibliographic sections of these different websites were manually navigated to look for

relevant documents. Indeed, due to time restraints, we restricted our manual research on specialist websites to available documents and reports only. Therefore, we did not include webpages or videos, which are often equally available. As it is commonplace that key method details are *too* summarised on webpages, we took the decision to only analyse ‘downloadable’ documents and reports enabling in-depth reading of texts.

Supplementary searches

A call for literature was conducted via two stakeholders: RNF and EuroParc. Specifically, a request for grey literature (scientific articles excluded) was undertaken through their mailing lists and professional networks. This allowed the mapping team to acquire 42 additional bibliographic documents (including technical reports, syntheses, and Master’s theses).

Testing the comprehensiveness of the search results

A test list of 40 scientific articles was used to assess the comprehensiveness of the search string on literature databases. The test list was composed of relevant scientific articles including a range of conservation translocation interventions, study populations, and publication years thought to be representative of the literature base and appropriate for answering the primary question. The review team identified these articles prior to the mapping process. Of the 40 articles 38 were retrieved by the search string (either in WOS, Google, or Scopus). Firstly, we checked if the search string retrieved articles in WOSCC. If an article was not retrieved by the search string in WOSCC, we checked for its retrieval in Google. If not retrieved in Google, we checked for its retrieval in Scopus. (If an article in question was indeed indexed in WOSCC, we did not check if indexed Scopus.) WOSCC retrieved 32 out of a total of 35 indexed articles. After checking for the remaining articles in Google and Scopus, an additional 2 were retrieved by Google, and 4 by Scopus (38 in total). Thus, the combined sub-comprehensiveness was 100% (38/38) if not including the 2 articles that were not indexed at all. But, the global comprehensiveness of our search strategy was 95% (38/40) as 2 articles not indexed in either WOSCC nor Scopus were not found either by Google. More details (including the two unretrieved articles) are provided in Additional file 3.

Article screening and study eligibility criteria

Screening process

All citations were extracted and archived in a bespoke excel spreadsheet. Screening and management of

bibliographic information was subsequently undertaken in excel. Three distinct screening stages were carried out successively for all articles (including articles coming from the call for literature and the specialist websites): (1) titles, (2) abstracts, (3) full-texts. Articles found by means of hand searching specialist websites were only assessed at the full-text stage. During the title and abstract screening process, in case of any doubt over the presence of a relevant inclusion criterion (or if the information was absent) the article was passed, by default, to the next screening stage.

The three screening stages were performed by two reviewers (JL and RS). In order to check that consistent and repeatable decisions were made, adherence to the eligibility criteria was assessed between the two reviewers using a Kappa test at the start of each screening stage. To that end, a set of articles was randomly selected: 10% for titles after duplicates were removed (463/4632), 10% of retained abstracts (193/1786), and 5% of retained full texts (59/1182). Kappa statistics were calculated at each stage, scores greater than 0.7 were deemed successful i.e. agreement attained. Any remaining disagreements were discussed and resolved between the two reviewers before commencing screening of literature. During the screening process, reviewers did not screen any articles that they had authored themselves.

Eligibility criteria

Article eligibility was based on the list of PICO criteria detailed in Table 2, with no deviation from the a priori systematic map protocol.

No study design types were excluded during the screening stages. This was done in order to achieve a comprehensive overview of the different kind of evidence existing on wildlife translocations in protected areas.

Regarding language, we included only publications written in English or in French. This criterion was considered at the full-text screening stage. This meant that if an article had an abstract written in a language other than French or English, it was still not excluded and it was transferred to the full-text screening stage.

No publication types (e.g. books, conference proceedings, peer-reviewed journal articles, theses, technical reports, etc.) nor contents (e.g. field studies, reviews, meta-analyses, discussion papers, etc.) were excluded during the screening process. However, reviews and syntheses (e.g. systematic reviews and maps, and meta-analyses) were not coded because these evidence syntheses often summarise findings across an overlapping set of multiple articles.

Study validity assessment

No critical appraisal was undertaken on retained articles because the intention of the map was not to examine the robustness of the study designs. Critical appraisals of study validity is not essential for systematic maps but we coded study design to provide a preliminary overview of internal validity [61].

Data coding strategy

Metadata were coded for all the articles passing the three screening stages. For this coding, the “study unit” was defined as “a unique translocation operation”. Thus, if multiple studies were reported within one article they were entered as an independent line in the systematic map database. In addition, one translocation operation was distinguished from another if the populations were different and/or if release site localities were unique geographical locations (cf. “Data mapping method”).

The key variables of interest were coded in terms of: (1) bibliographic information (the document type, and type of bibliographic content), (2) study characteristics (the study country, release and capture coordinates, climate zones of capture and release sites, distance (km) between each, source and destination e.g. captive site to wild release, protected area context and IUCN protection category, the release site’s biome, and programme’s motive, period of time, costs), (3) population characteristics (the taxonomic group, sample size, age of released individuals), (4) intervention characteristics (whether a reintroduction, supplementation, introduction, or a combination of two), and (5) outcome characteristics (whether related to physiology, feeding, behaviour, space use, reproduction, survival, demography, or genetics). Full details are given in the systematic map protocol, with no deviations (a codebook is provided along with example coding formats). When the data was not sufficiently detailed or simply unknown, we coded as such (using “Unknown” when necessary).

The coding stage was performed by three coders (JL, RS and HM) who shared the eligible articles between them and coded all variables for their respective samples. Due to resource and time limitations, an independent dual coding of the entire database was indeed not feasible. However, in order to guarantee that information was coded and extracted in a consistent and repeatable way, a four-step process was undertaken.

- Step 1 “discussion session”: discussions were held between the three coders, before the extraction of metadata commenced, to ensure any doubts prior to coding were clarified: JL gave thorough explana-

tions regarding coding rules to RS and HM for all the characteristics of the Map.

- Step 2 “practice and re-discussion session”: immediately following the first discussion session, 5 articles were randomly selected to be practice coded by RS and HM. This coding step was monitored by JL who gave further explanations to the two other coders in cases of doubt. Indeed, any disagreements were re-discussed. This was a pre-pilot testing phase, purely for practice purposes.
- Step 3 “pilot-testing”: a pilot extraction phase was undertaken by JL, RS, and HM on an identical sample of 10 articles. This was undertaken to ensure all characteristics were coded consistently by all three reviewers, discussing any potential disagreements. Before starting the actual coding process, this pilot-testing step was repeated until no disagreements occurred. We required two tries until all coded characteristics were homogenous. 20 articles were therefore coded in total.
- Step 4 “post-coding crosscheck”: once coding had been completed, an a posteriori crosscheck was carried out by JL on the entire database to identify any errors and homogenize all coded terminology. All disagreements were discussed until a consensus was reached for the concerned articles. (*N.B.*, during the coding process the reviewers were in contact and could question each other in case of any doubt).

Data mapping method

All the coded metadata for all the studies were included in a systematic map database: a bespoke

Microsoft excel spreadsheet. By cross-tabulating the data, summary figures and tables were produced for this Map report to identify knowledge gaps (i.e., subtopics requiring further primary research) and knowledge clusters (i.e., subtopics that are sufficiently covered by existing studies to allow for full systematic reviewing). The distribution and frequency of translocation operations was presented with heatmaps, or geographical maps illustrating the number of interventions by country. Based on these results, recommendations were made on priorities for policy makers, practitioners, and research.

Because our aim was to not only code the literature base in terms of number of publications (hereafter referred to as “Bibliometric synthesis”) but also extract information on specific translocation events within these retained publications (hereafter referred to as “Translocation synthesis”), we underwent a three-step mapping process:

- Firstly, we permitted a coding of “unclear context” due to the complexity of many publications. If publications did not explicitly describe the name of a protected area in the given translocation event but gave sufficient information to assume our PICO criteria were otherwise respected, during the screening process, these articles were coded but consequently not included in the systematic database. A brief narrative summary is given without tables (cf. “Unclear cases” section, in [Review findings](#)).

Table 2 Inclusion/exclusion criteria for the three-stage screening process

Eligible population(s)	All terrestrial, marine, or aquatic plant or animal populations from any country across the globe that were of wild or captive source. Wild populations included any free-ranging species from natural environments; not domesticated or cultivated. Captive populations included species born in captive setting (e.g. zoos, nurseries, or breeding pens)
Eligible intervention(s)	All conservation-based translocation interventions occurring either (1) between two different protected areas; (2) from a non-protected area to a protected area; (3) from a protected area to a non-protected; or (4) within the same protected area where the aim is improving the viability and persistence of the translocated population (i.e. reintroduction, supplementation, or assisted migration (see Table 1 for definitions). Ecological replacement was also included as an eligible intervention where the goal is to re-establish an ecological function lost through extinction, which involved the translocation of the most suitable existing sub-species, or a close relative of the extinct species
Eligible comparator(s)	With regard to a systematic map, comparator was not a screening criterion
Eligible outcome(s)	Any outcome-related effects on the translocated populations. This included (1) Space use i.e. studies measuring all movement/dispersal of translocated individuals, notably home range measurements or Euclidean distance travelled. (2) Demography i.e. studies outlining the changes in number of individuals (males and/or females) of the translocated population. (3) Survival i.e. studies illustrating precisely the proportion of individuals alive or level of mortality since translocation. (4) Reproduction i.e. number of young born since translocation, or specifically the survival rate of offspring. (5) Feeding i.e. all effects specifically on diet and feeding of translocated individuals. (6) Behaviour i.e. studies measuring changes in terms of communication (e.g. vocal), social structure, or anti-predator behaviour e.g. stress/vigilance levels, of translocated individuals. (7) Genetics i.e. studies relating to the genetic structure of the translocated species. Finally, (8) Physiology i.e. biological or physiological impacts measured at the molecular, cellular or organic level of translocated population (e.g. hormone activity)

We divided the Map's narrative synthesis into two parts, a "Bibliometric synthesis" and a "Translocation synthesis".

- ii. Concerning the Bibliometric synthesis: this part of the synthesis included all retained literature coded by document type i.e. one line in the Excel datasheet corresponding to a single publication (i.e. either a Book, conference object, journal article, report, technical documentation, or thesis). For each document type, we coded bibliographic content (i.e. either a discussion paper, modelling, overview, review, or study). *Study* in this instance was regarded as a *field-based* study where data was acquired in the field. Equally, "programme motivation" was coded except for reviews and overviews because these documents may themselves summarise a number of operations with a number of different motives. Thus, reviews and "metapopulation management" (one of six coded programme motivations) articles were coded only for the bibliometric variables mentioned above. Concerning the latter, these were considered inappropriate and too time consuming to code in the translocation synthesis as the same individuals may be repeatedly transferred back-and-forth between fenced reserves in order to cope with the particular problem of lack of gene flow between enclosed populations.
- iii. Concerning the "Translocation synthesis": from all the remaining publications this part of the synthesis involved coding the specific translocation events consistent with the IUCN definitions. Two key variables (method and location) were used as cut-offs to define a study unit [63], i.e. a translocation. Therefore, a translocation was coded as a unique data line in the Excel datasheet if: (1) a different species was translocated, and/or (2) the release site was a unique geographical location. If the same species were translocated to clearly different release sites, these were presumed not to be the same individuals under movement thus coded as unique data entries. Time duration was not considered as a cut-off when distinguishing translocations events. Consequently, some translocation interventions, coded as for example, "Reintroduction + supplementation" may have particularly long time periods if the supplementation part of the intervention followed the initial reintroduction several years later.
- iv. From this, a search for linked data (cf. "Linked data" section) eliminated duplicated translocations from the evidence base.

Linked data

Once all the studies had been coded, we undertook a search for linked data. This was done in order to locate articles, published by either the same or different authors, referring to the same translocation event. Indeed, we wanted our Map to be able to isolate single translocation events, without duplicates i.e. multiple publications referring to the same event. Therefore, if: (i) the same species, (ii) the same capture site, (iii) the same release site, (iv) the same intervention type, (v) the same number of individuals reported, and (vi) for the same translocation period, we subsequently tagged such translocations as "linked data". We assumed that if these 6 factors matched, the likelihood was high that two different publications were reporting the same transfer of individuals. When a single translocation was coded multiple times, we retained the publication providing the more detailed description. In the case where a single event was described in a thesis and a corresponding scientific article, we retained the peer-reviewed article.

Review findings

Literature searches and screening process

The ROSES flow diagram for systematic maps illustrates the screening process (Fig. 1). Our literature search yielded 4981 hits from WOSCC and Scopus. A further 1373 hits from various other sources were retrieved: Google Scholar, organisational websites, a grey literature call and hand search on specialist websites. This resulted in a total of 6354 records and 4617 unique records after removal of duplicates. At title screening, 2841 titles were excluded, leaving 1776 for abstract screening. During the abstract screening stage, 510 references were excluded, leaving 1266 for full-text screening. However, 78 articles (6.2%) were unobtainable (Additional file 4) i.e. authors were not able to locate the articles in question through the internet or French National history Museum or the CNRS (National Scientific Research Centre) subscriptions. Thus, 1188 articles were submitted to full-text screening. A total of 498 articles were rejected at full-text mostly due to irrelevant interventions (i.e., articles in question did not describe a translocation operation in the context of a protected areas), irrelevant outcomes (i.e., article did not evaluate effects on translocated population), or irrelevant population (i.e., article reported data for invasive species, or species manually relocated for non-conservation purposes such as hunting). Finally, a total of 690 publications were accepted for coding [Additional file 6 sheet "Full List of articles (690)"]. Additional file 5 outlines all information on the three screening stages

(inclusion/exclusion screening process, dates, screeners, reasons for exclusion, etc.) (sheet “List of all articles”). The list of articles excluded at full-text stage along with reasons for their exclusion can be found more easily in a specific sheet of the Additional file 5 (sheet “List of excluded articles FT”).

From the 690 articles accepted as eligible for the review after the three screening stages,

- i. 42 publications (65 translocations) were coded as “unclear context” not sufficiently describing the role of the protected area in the given translocation event despite these articles meeting our PICO during the screening process. These articles were consequently not included in the systematic database but in a specific Additional file 6 (sheet “Unclear contexts”) and described briefly in “Unclear cases” section in [Review findings](#).
- ii. 58 linked translocations (35 publications) were detected and eliminated from both parts of the narrative synthesis (see Additional file 7, sheet “Linked data”).
- iii. Concerning the Bibliometric synthesis [Additional file 7—sheet “4. Full database (613)”]: 613 publications in all were coded for the following bibliometric variables: document type, bibliographic content, and programme motivation). Except programme motivation was not coded for 90 reviews (cf. “[Data mapping method](#)”).
- iv. Concerning the “Translocation synthesis”: 841 translocations (i.e., intentional movements of any organism from one site to another) were fully coded in respect to all descriptors [see Additional file 7, sheet “Translocation database (498)”. These translocation events were extracted from 498 different publications (having not extracted translocation events from 115 publications: 90 review articles and 25 “metapopulation management” articles).

Bibliometric synthesis

The total number of publications summarised in this section is 613—after eliminating “unclear contexts” and “linked data”—including reviews, programme motivation, and translocation studies. (cf. sheet 4 “Full database (613)” in Additional file 7).

Publication type and content

Figure 2 shows the distribution of publication types. The systematic map is mainly composed of journal articles (85%). The second highest proportion (5%) of publication type in the systematic map are book chapters.

Ph.D. theses accounted for 3% of included publications. This equated to 15 different theses reporting translocations in the context of protected areas. Figure 3, shows the distribution of publication contents. Indeed, the majority of translocations in the contexts of protected areas were published from field studies i.e. data acquired from on-the-ground studies. Then, reviews (9%), and then other (8%, comprising of various publication formats such as short communications). Overviews made up 5% of the systematic map, followed by modelling publications (3%) where field data was not necessarily collected.

Publication motivation

In terms of programme motivation by publication (Fig. 4)—not including the 90 review papers—the vast majority of articles published documented aims to restore and improve the conservation status of the focal species: 353 publications in total, which represented 57% of all publications. Secondly, 9% (56) of publications did not detail the motivations for translocating wildlife (“Unknown”). 8% of publications were for documenting a trial transfer prior to a translocation operation. 25 publications were specifically on metapopulation management, 18 on wild-human conflicts, 18 on wildlife rescue, with only a very small number of articles on rewilding (1%).

Publications by year


Publication dates ranged from 1969 to 2020, with the majority after 1998 (93%) (Fig. 5). The period 1969 to 1997 equated to 7% of publications. Grey literature made up a larger proportion of the total studies in more recent years. Indeed, the grey literature included in the systematic map were all published from 1994 onwards.

Translocation synthesis

The number of publications summarised is 498, from which 841 unique translocations were coded. (cf. sheet 5 “Translocation synthesis” in Additional file 7). 115 publications (Reviews and “metapopulation management” papers) are excluded from the translocation synthesis section.

Translocation motivation: reasons for moving wildlife

We coded 622 translocations, equating to 78% of all coded interventions in the evidence base as undertaken in order to improve a focal species’ status. Second to conservation purposes, were wildlife rescue operations, this equated to 58 of all translocations carried out in the context of protected areas. Surprisingly 56 translocations comprised of trial or experimental transfers of wildlife to protected areas pre-dating conservation



programmes. Human-wildlife rescue operations accounted for 33 translocations, unsurprisingly only animals are concerned, of which 17 transfers did not document if individuals were released into resident populations or not (Table 3). Regarding rewilding, which was considered in the context of reintroducing a species to restore lost ecological functions, equated to only 8 translocations (4 in Argentina, 2 in Brazil and 2

in Germany), and in each case the relocation of a mammal. One study, in Brazil, aimed at reintroducing the red-humped agouti and the brown howler monkey, in Tijuca National Park, Rio de Janeiro; two species that are important seed dispersers, which may be particularly effective at restoring such ecological processes [64]. Finally, 64 translocations had completely

unknown motives, even if their intervention type was detailed (Table 3).

Translocations by year

In terms of actual translocations by year (Fig. 6), we can observe a similar trend to that of Fig. 5. Over time, the number of translocation operations have accelerated. Overall, from 1997 to 2020, on average $22.6(\pm 20.9)$ translocations (all combined) per year were undertaken. But since 2018, an average of 54 per year have been published. From the current evidence base, the year with the highest number of published translocations was that of 2018 (85/841). Further still, only 6% of coded translocations were undertaken before 1997. This increase in research output may be an indirect response to the increasing pressures currently facing many animal and plant populations [65], in that in the number of performed conservation translocations are increasing because it is seen as a viable way of enhancing the resilience of threatened species [25, 27, 66], and thus publications have increased by consequence.


Study locations and climate zones

Translocations included in the systematic map spanned the globe (Fig. 7). Most translocations were performed in North America, Oceania, and Europe. There was a clear lack of information on translocations in Northern Africa and the Sahel region, and Central Asia. Of the 68 included countries, the most translocations were

undertaken in the USA (22%), Australia (16%), New Zealand (10%), and Canada (4%). In accordance with this geographical distribution, the majority of translocations were released in oceanic/maritime climates (Cfb: 234), followed by warm temperate climate (Cfa: 84), Tropical savanna climate (Aw: 65), humid continental climates (Dfb: 59), desert or arid climate (BWh: 50), equatorial climate (Af: 40), and semi-arid climates (BSk: 37) (Figs. 8 and 9). Less than half of translocations gave sufficient information in order to code the capture and release site climates (44%). Of the known climates zones (358 translocations), 11% of translocations changed climate zones between capture and release sites.

Study population: which taxonomic groups are translocated?

Translocations that were included in the systematic map mainly involved animal species (686). Although, 148 translocations of plants were also included, and 7 translocations of fungi. Of the animal translocations, mammals were the most translocated group (Fig. 10), almost 400 (equating to 56%) of reported translocations between 1969 and 2020 from 45 different taxonomic families, and 158 different species (Additional file 7). Over 170 (25%) translocations involved birds, followed by reptiles (42), fishes (28), insects (21), and amphibians (15 translocations). From mammals, the most commonly translocated taxonomic orders were






Fig. 3 Proportion of publication contents

Artiodactyla (of which *Bovidae*, and *Cervidae* were the most translocated families), *Carnivora* (of which *Canidae* and *Felidae* were the most translocated), *Perissodactyla* (of which *Equidae* were most transferred), *Diprotodontia* (of which *Macropodidae*), Primates (of which *Hominidae*), and *Rodentia* (of which *Castoridae* was the most relocated). Although marine fauna are also catalogued—*Bivalvia*, *Anthozoa*, and *Hexanauplia*—it would appear that they receive much less attention (Fig. 10). Several plant groups were also translocated (Fig. 11). However, a large bias towards *Magnoliopsida* is evident, followed by *Liliopsida*, and *Lecanoromycetes*. Among translocated *Magnoliopsida*, *Asteraceae* were by far the most translocated family. However, a distinct lack of literature concerning non-vascular plants such as bryophytes is highlighted. These knowledge gaps are likely to be associated with the complexity of transferring such taxa. The number of individuals translocated, by age class and taxonomic groups, are provided in Table 4 for *Animalia* and Table 5 for *Flora*, for all studies and in average (\pm standard deviation).

Study interventions: what type of translocations exist, how many individuals are translocated, to where, and over what distances?

Tables 6 and 7 illustrate the distribution of the number of translocation operations by intervention type, taxonomic class and age at release, for animals and plants/fungi respectively. Concerning animal translocations (Table 6), 686 were coded in total. 12 different taxonomic classes (332 species) were subject to conservation translocations. The most common intervention was that of “reintroduction + supplementation” (176) i.e. a founder group of a species previously extinct or extirpated reintroduced into its historical range and later reinforced. 158 translocations were of one-off reintroductions (the founder group not being reinforced at a later date) and supplementations. Note that 182 interventions were coded as “Unknown” as information was lacking in their methods. The most translocated taxonomic group were mammals and birds. Of these translocations, adults only or a combination of adults and juveniles was most often the chosen strategy. For instance, adults were transferred alone in 104




Fig. 4 Illustrating the 6 main motivating reasons for translocations. This combines all translocations interventions (introductions, reintroductions, and supplementations). Figures describe motivation by number of retained publications (thus “no coding” refers to the 90 review publications where metadata for motivation was not extracted, and “metapopulation management” refers to 25 publications where individual translocation were not extracted)




Fig. 5 Number of translocation-related publications per year

Table 3 Heat-map showing the distribution of the number of translocation operations in terms of programme motivation and taxonomic kingdom. Conservation = any attempt to restore or improve the status of the translocated species; Experimental or trial = a trial relocation prior to the main event; Human-wildlife conflict = any interaction between humans and fauna resulting in the relocation of the individual(s) in question; Rewilding = reintroduction or introduction to restore a lost ecological function; Wildlife rescue = a relocation in order to save the individuals from human activity; Unknown = no information (or not detailed enough) provided by the authors to class motive

Taxonomic kingdom X Programme motivation	Intervention type						Total
	Intro+suppl	Introduction	Reintro+suppl	Reintroduction	Supplementation	Unknown	
Animalia	6	6	176	158	158	182	686
Conservation (improving status of focal species)	6	4	158	123	110	88	489
Experimental or trial translocations		1	4	12	13	16	46
Human-wildlife conflict				5	11	17	33
Rewilding (restoring natural functions)			3	3		2	8
Unknown		1	9	11	9	33	63
Wildlife rescue operation			2	4	15	26	47
Fungi				4	3		7
Wildlife rescue operation				4	3		7
Plantae		4	10	11	41	82	148
Conservation (improving status of focal species)		3	10	9	39	72	133
Experimental or trial translocations		1		2	2	5	10
Unknown						1	1
Wildlife rescue operation						4	4
Total	6	10	186	173	202	264	841

“Intro + suppl”: the movement of individuals to a habitat outside of their historical range, and subsequently followed up by a supplementation event

Introduction: a one-off transfer of individuals to a habitat outside of their historical range

“Reintro + suppl”: the movement of individuals that were otherwise extinct or extirpated to a habitat within their known historical range, and subsequently followed-up by a supplementation event

Reintroduction: a one-off transfer of individuals that were otherwise extinct or extirpated to a habitat within their historical range

Supplementation: a one-off transfer of individuals to reinforce an existing resident population

Unknown: refers to all movements that were not detailed sufficiently by the authors in order to code the type of intervention with high confidence

N.B., white space equates to no available data. Red colours equate to a few number of translocations, green colours represent a greater number





Fig. 6 Number of translocations per year




mammal translocation operations vs. 10 translocations where only juveniles were transferred. The least translocated groups were marine taxa, the logistical complexity of such interventions and the specialised equipment such as protective cages [69] needed to increase the chances of success likely, to some extent, explain the lack of literature.

Concerning translocation of plants or fungi (Table 7), 155 translocations, spanning at least 5 different taxonomic groups and 1 fungi taxon in total were documented. The most common intervention type was “supplementation” (44). However, similarly to animal translocations, a large number were uncodable interventions arose due to a lack of information in methods (82). A diversity of plant development stages were translocated. However, contrary to animal translocation, a single development stage (i.e. ‘age group’) was most commonly selected for translocation. For instance, concerning *Magnoliopsida*, only one translocation operation attempted to transfer adult plants,

seedlings and seeds vs. 23 operations which chose to transfer seeds only.

Further, in terms of the number of individuals transferred per translocation, Tables 4 and 5 summarise this profuse information across the years (1969–2020) in terms of age by taxonomic groups, respectively for animals and plants/fungi. Indeed, concerning animal translocations, over time the number of individuals involved in mammal translocations has been far higher than other taxonomic groups (>71,000 individuals translocated since 1969). Indeed, the average number of individuals transferred per translocation was 529, with a median number of individuals transferred per translocation equating to 20. Interestingly, concerning amphibians, few translocations (15) have been carried out, but large numbers of individuals (17,801) were used across operations. Indeed, it is often the case that large numbers of amphibians are needed to ensure that a viable proportion of the founder population persists [70].




Regarding plant translocations, the highest number of individuals per translocation operation was highest for *Magnoliopsida* sp. followed by *Pinopsida*, and then *Liliopsida*. Interestingly 7 translocations attempted the transfer of thallus fragments of fungi groups (*Lecanoromycetes*). Concerning *Magnoliopsida* translocations, the median number of seeds transferred per translocation was >3000, where as if adult plants only were chosen for transfer, the median number was 35 per translocation.

In terms of direction of transfer (Fig. 12), the majority (592 translocations) involved a transfer of individuals from non-protected sites to protected sites (illustrated by the “To” category, in Fig. 12). However, of these transfers, 64 had unknown source sites. Secondly, 196 transfers of species were from one protected area to another (“From-To”), 47 translocations occurred within the same protected area (“Within same”), and in 27 rare cases, species were translocated from a protected area and release into non protected sites (“From”).

Concerning translocated animals only from non-protected to protected sites (“To”), 204 of these translocations included the procurement of individuals from either captive or breeding sites subsequently released into protected wild areas. 209 involved the transfer of animals from wild sites to another wild site. Furthermore, these translocations had diverse motives from rescue operations and human-wildlife conflicts, to pure conservation purposes. In two specific examples, a conservation-motivated reintroduction of *Ursus arctos*, where ten adults were transferred from the “Slovenian alps” to Adamello Brenta Nature Park in Italy [71] was coded as wild site to wild site. In another example of a conservation-orientated reintroduction, 43 passerine birds were transferred from a wild site (verified by given coordinates) to Mulligans Flat Nature Reserve in Australia [72].

Concerning habitat protection, of animal translocations, 20 (13 mammals, 4 insects, 3 birds) were known to be translocated to a protected area with a lower IUCN protected status than the source site, according



to Protected Planet (<https://www.protectedplanet.net/en>). In one instance [73], this involved the relocation of 201 *Cervus canadensis* from Elk island, Alberta (IUCN protection status: II) to a protected landscape (V) over 4000 km away: the “Tennessee Cumberland mountains”. In another [74], 687 *Deinacrida mahoenui* were transferred from an “scientific reserve” (Ia) to a “scenic reserve” (III), in New Zealand. Comparatively, 38 species were moved to a protected area with a higher protection status. 62 translocations did not experience change in protection status. For 567 translocations of animal species, the protection status’ were not reported thus protection status change was *Unknown*. Concerning the 155 plants and fungi translocations, very little information was reported in respect to protection status: six translocations where no protection status

change occurred. The rest of the events were coded as *Unknown*.


Concerning distances, the map team were able to code up to 358 translocations where geographic coordinates were retrievable for both capture and release sites (representing approximately a third of the “translocation synthesis”). Additional file 8 shows the methods used to calculate distances for these translocations. Indeed, we used the standard Mercator projection. A diverse number of distances are evident for animal translocations. Indeed, the majority of cases were short distance translocations i.e. 0–100 km. However, we can also observe that 54 animal translocations carried out involved distances of over a 1000 km (Fig. 13). Concerning plants, only 3 translocations involved distances greater than 100 km.

Study outcomes: which impacts are measured?

Reported outcome measures largely concentrated on: space use, survival, demography, reproduction, feeding, behavior, genetics, and physiology (Fig. 14). Outcome metrics related mostly to the survival of individuals translocated (367). Secondly, space use was largely studied in order to determine all movement/dispersal of translocated individuals. This included notably home range measurements, or straight-line distance travelled for animals. There were also a number of studies that focused on demography metrics (220 translocations) i.e. changes in the number of individuals, males/females, of the translocated population overtime. Reproduction was the fourth most studied outcome metric (165 translocation studied reproduction in the following months or years). This included any study expressing findings on the number of young born since translocation. Comparatively, few studies focused on physiology (72), behaviour (83), genetics (68), and feeding (57). These are still important aspects to take into account especially when attempting to define if a project is a success or not. In the case of behaviour, social structure and cohesion, social roles and social learning may all be important to consider in building populations that are resilient and likely to persist in the long term [75]. Also, understanding the physiology and wellbeing of individuals either before or after release are also important factors to consider [76], and may be particularly useful in predicting long term survival rates for future interventions.

Financial costs

Financial costs were reported in only 23 translocations. Large differences in costs were reported. This was notably due to a choice of expenditure details expressed by authors. Some articles reported total project costs,



while others, for instance, only reported transfer modality costs excluding salaries or other expenditures commonly associated with translocation programmes. For example, Kenup et al. [77], reported total costs of \$6300 which included monitoring, captivity and release of *Dasyprocta leporina*. Whereas Bennett et al. [78] reported total expenditures of up to \$185,000 for a reintroduction project of *Climacteris picumnus*. Baker et al. 2012 supplementation efforts including 50 individuals of *Dasyornis brachypterus* cost upwards of AU\$200,000 [79], which included total expenditures from financing Ph.D. students to capture and release modalities. Muths et al. [80], an intervention including the transfer of eggs (*Bufo boreas*), equated to \$700 (excluding salaries).

A guide for managers

Finally, due to the complexity, and quantity of data available in this map, we realise that not all data was able to be summarised. Thus, we provide a ‘guide’ for managers to navigate through the evidence base of Additional file 7 (Fig. 15), in order to gather data that they may perceive necessary and access the references of the original studies utilised for any aspect of the map that is of interest. Among other information, this should allow users of the map to extract information on the factors of success. However, attention must be applied to the term “success”; we have not attempted to code if a study was successful or not because from a biological point of view this depends on many factors. Hence, we have only coded the factors that affect success (e.g. number of individuals translocated, the

quality of release habitat which is indirectly illustrated by the IUCN protection status, the distances involved, whether acclimatisation measures are used). The evidence base provides an overview of existing management interventions, summarised in association with such important factors that must be considered when engaging in translocations.

Unclear cases

Our mapping methods were conservative in nature, this meant that a total of 65 translocations (39 animals, 26 plants) were excluded from the descriptive statistics due to lack of sufficient confirming evidence specifically on protected area context. However, there was also a lack of confirming information to outright exclude such publications during the full text screening stage. Authors conclude that these instances probably should, in truth, have been included in the map description but did not want to risk potential ‘error bias’. Indeed, the authors of the map team feel that these cases merited coding for the rest of the variables and inclusion into an additional file (cf. sheet 5 of Additional file 7). Of these 65 translocations: two were introductions of animals (one bird, one fish species); 5 were one-off reintroductions (2 species of *Mammalia*, 1 species each of *Aves*, *Mollusca*, and *Magnoliopsida*); 11 reintroductions followed by supplementing the same species (concerning animals: 3 mammals, 1 fish, 1 bird, 1 amphibian, 1 *Bivalvia*. Plants: 3 *Magnoliopsida*, and 1 *Liliopsida*). 36 interventions were coded as *Unknown* due to lack of confirming information on intervention type.

Table 4 Illustrating the total sum of individuals (*Animalia*) translocated across all years (1969–2020) by taxonomic class

Animal taxonomic class	Age class as stated by authors							Total ind translocated	Total number of coded translocation operations
	Adults only	Juveniles only	Adult and Juveniles	Adult, Juveniles, and Larva	Instar only	Larva only	Unknown age class		
Mammalia	55,094 (529 ± 5129) 20	310 (31 ± 61) 38	6147 (81 ± 165) 51	–	–	–	10,159 (64 ± 127) 29	71,710	383
Aves	910 (43 ± 35) 18	4637 (122 ± 160) 46	1327 (60 ± 39) 56	–	–	–	6160 (87 ± 189) 29	13,034	175
Reptilia	296 (27 ± 16) 23	149 (75 ± 100) 38	1081 (77 ± 82) 51	–	–	–	4684 (335 ± 673) 65	6210	42
Actinopterygii	–	1955 ^a	6516 (2172 ± 1811) 69	–	–	–	6420 (279 ± 550) 30	14,891	28
Insecta	5995 (545 ± 983) 20	644 (161 ± 57) 36	–	–	1010 ^a	220 ^a	1350 (337 ± 261) 26	9219	21
Amphibia	6031 (2010 ± 3378) 21	12,424 (3106 ± 5264) 38	1594 ^a	3783 ^a	–	–	–	23,832	15
Anthozoa	–	–	–	–	–	–	65 (5 ± 6) 22	65	13
Gastropoda	–	–	765 (383 ± 388) 45	–	–	–	192 (96 ± 20) 31	957	4
Bivalvia	–	–	–	–	–	–	200 (100 ± 71) 27	200	2
Athropoda	–	–	–	–	–	–	60 ^a	60	1
Hexanauplia	–	–	–	–	–	–	–	Unknown	1
Malacosraca	62 ^a	–	–	–	–	–	–	62	1
Total ind translocated	68,388	20,119	17,430	3783	1010	220	29,290	140,240	686


In parentheses: average ± standard deviation. Figures in bold represent the median number of individuals translocated. '–' no available data. "Unknown" signifies that for the given translocation events exact numbers of individuals are not known

^a Denotes a single translocation event, thus no median, mean or standard deviation are calculated

Comparison to other reviews

Relating our current findings to others evidence syntheses may be useful for managers and policy makers. To our knowledge no other systematic map exists on wildlife translocations. However, map-like reviews (hereafter referred to as 'overviews') do exist providing some state of knowledge. But given the large differences in terms of methodology, topic scope, and volume of literature exposed in Table 8, it was deemed difficult to make legitimate graphical comparisons. For instance, of the 4 overviews compared to our map, none included all biodiversity (terrestrial and marine animals, plants and fungi), or else their exclusion criteria was difficult to ascertain. Concerning three overviews of which only animals were the scope, mammals and birds were consistently the most translocated groups. Fischer and Lindenmayer [45], found that over

50% of studies considered were performed with mammals and approximately 43% with birds. In an overview on key-stone species, 54% of publications focused solely on mammals [55]. Further, in a recent overview by Rescendre et al. [56], *Diprotodontia* was the highest represented mammal order, although in our map it was *Artiodactyla* (notably, but not exclusively, *Bovidae*, and *Cervidae*). Concerning supplementations, Fischer and Lindenmayer [45] showed that only birds and mammals were involved, although our map synthesises information from 8 different animal taxonomic classes, 3 plant classes and 1 fungi class in respect to supplementations. In terms of release strategies, birds were most commonly soft released [51, 56], our map also adheres to their findings: 67/175 bird translocations used soft release measures. Although in our map, release measures for 95/175 were unknown. Despite the diversity of



methods, we are able to demonstrate some consistency with previous overviews. However, the differences justify the need to have undertaken a systematic search of the existing literature in order to provide a clearer and more comprehensive state of knowledge.

Limitations of the systematic map

Search string

The search strategy used to generate this systematic map was designed to capture a large breadth of relevant literature, not only on wildlife translocations but specifically in the context of protected areas. However, there does exist a certain diversity of terminology used on the subject of translocations (i.e., some terms may have been missed that could result in bias in our evidence map). For example, in general there were fewer studies evaluating plant translocations. In our search string, we did not use specific terms such as transplantation or transplant* which may have otherwise retrieved more studies on plant species. However, we did carefully scope our search string for intervention terms by including several synonyms (e.g. “assisted colonization”, “assisted migration”, etc.). In addition, our comprehensiveness was very high (see “Testing the comprehensiveness of the search results” section).

Language

Translocation is often carried out by managers of natural areas or communities. It is therefore possible that the reintroduction experiences are largely recounted in grey literature because these actors are less used to producing academic literature. Also, this type of document

is often published in the language of each country rather than in English like scientific articles. Our search strategy included several sources of grey literature (4 web-based search engines, 5 specialist websites and an explicit call for grey literature). However, the fact that only English terms were used for our searches and that the call for literature was broadcast in French may have limited the recovery of grey literature written in a native languages other than English and French. In addition, 28 articles retrieved by our English search terms were not written in English and were excluded during the full-text screening stage in accordance with our screening criteria (Additional file 9): Spanish, 8; German, 7; Chinese 5; other languages, 8. However, it is unclear how many of these articles would have met all the inclusion criteria. Finally, our systematic map is likely to have underestimated the reality of the translocation cases regarding these language limitations due to a lack of time and skills.

Consistency

As explained in “Deviation from the protocol” section, we were not able to perform our Kappa test on 10% of retained eligible full-texts. Due to time constraints, we based our kappa test on 5% of the eligible corpus. We are aware that the assessment of our consistency at the full-text screening stage is therefore necessarily lower than expected. Nevertheless, 5% of all the full-texts to be screened (1182) represented 59 articles; this is already a large enough sample to meet different types of publications and to highlight possible disagreements. Moreover, the result of this full-text Kappa test

Table 5 Illustrating the total sum of individuals (*plantae and Fungi*) translocated across all years (1969–2020) by taxonomic class

Plant taxonomic class	Age class as stated by authors												Total ind translocated	Total number of coded translocation operations
	Adult plants only	Adult plants and cuttings	Adult plants, seedling, and seeds	Bulbs only	Cuttings only	Plantlets only	Rhizomes only	Seeds only	Seeds and seedlings	Seedlings only	Thallus fragments	Unknown		
Magnoliopsida	1090 (68 ± 62) 35	130 ^a	–	–	570 ^a	300 ^a	–	6750 (3375 ± 2015) 3375	480 ^a	45,912 (534 ± 1697) 106	–	6 (3 ± 3) 5	55,238	125
Liliopsida	340 (68 ± 97) 35	–	–	Unknown	–	–	–	–	–	2170 (271 ± 233) 106	–	10 (5 ± 0) 5	2520	16
Lecanoromycetes	–	–	–	–	–	–	–	–	–	–	475 (68 ± 60) 25	–	475	7
Syntaxa	–	–	–	–	–	–	–	–	–	–	–	–	Unknown	3
Pinopsida	–	–	–	–	–	–	–	–	–	2720 (1360 ± 905) 106	–	–	2720	2
Bryophyta	–	–	–	–	–	–	–	–	–	–	–	–	Unknown	1
Lycopodiopsida	20 ^a	–	–	–	–	–	–	–	–	–	–	–	20	1
Total ind translocated	1450	130	–	–	570	300	–	6750	480	50,802	475	16	60,973	155

In parentheses: average ± standard deviation. Figures in bold represent median number of individuals translocated

‘–’ no coded figures. ‘Unknown’ signifies that for the given translocation events exact numbers of individuals are not known

^a Denotes a single translocation event, thus no median, mean or standard deviation are calculated

Table 6 Number of translocation events (whether reintroduction, supplementation or introduction or a combination) for animals, by taxonomic class and age at release

Taxonomic class X age	Intervention type.					
	Intro+suppl	Introduction	Reintro+suppl	Reintroduction	Supplementation	Unknown
Actinopterygii			4	2	10	12
Adults + Juveniles			3			
Juveniles only			1			
Unknown				2	10	12
Amphibia	1		2	3	2	7
Adults only				1	1	1
Adults + Juveniles						1
Adults + Juveniles + Larva			1			
Eggs only						1
Juveniles only	1		1	1	1	
Unknown						5
Anthozoa					13	
Unknown					13	
Athropoda						1
Unknown						1
Aves	2	2	54	43	40	34
Adults only		2	6	8	3	2
Adults + Juveniles			11	2	6	3
Juveniles only	1		12	6	13	6
Unknown	1		25	27	18	23
Bivalvia						2
Unknown						2
Gastropoda				1	3	
Adults + Juvenile					2	
Unknown				1	1	
Hexanauplia						1
Unknown						1
Insecta	1	2	5	4	4	5
Adults only	1	2	2	1	3	2
Instar				1		
Juveniles only			1	1	1	1
Larva only			1			
Unknown			1	1		2
Malacostraca				1		
Adults only				1		
Mammalia	2	2	101	94	75	109
Adults only		1	23	19	32	29
Adults + Juveniles			25	13	17	22
Juveniles only			1	3	5	1
Unknown	2	1	52	59	21	57
Reptilia			10	10	11	11
Adults only			5	2	3	1
Adults + Juveniles			4	4		6
Juveniles only					2	1
Unknown			1	4	6	3

Table 7 Number of translocation events (whether reintroduction, supplementation or introduction or a combination) for plants and fungi, by taxonomic class and age at release

Taxonomic class X age	Intervention type				
	Introduction	Reintro+Suppl.	Reintroduction	Supplementation	Unknown
Fungi			4	3	
<i>Lecanoromycetes</i>			4	3	
Thallus fragments			4	3	
Plantae	4	10	11	41	82
<i>Bryophyta</i>			1		
Adult plant			1		
<i>Liliopsida</i>		3	1	9	3
Adult plants only				4	1
Bulbs only		1			
Seedlings only		2	1	3	2
Unknown				2	
<i>Lycopodiopsida</i>			1		
Adult plants			1		
<i>Magnoliopsida</i>	4	7	8	31	75
Adult plants only			5	5	6
Adult plants + Cutting			1		
Adult Plants + Seedling + Seed		1			
Cuttings only				2	1
Plantlets only		3			1
Rhizome only		1			
Seeds only	1				1
Seeds + Seedlings				1	
Seedlings only	1	2	1	23	63
Unknown	2		1		3
<i>Pinopsida</i>				1	1
Seedlings only				1	1
<i>Syntaxa</i>					3
Seeds only					3

"Syntaxa" refers to where more than one class were translocated at the same time

was excellent (0.93) with disagreements only on two full-texts. This confirmed that carrying out the full-text Kappa test on 5% was not detrimental to the rigour of the systematic map.

Regarding the coding step (extraction of the meta-data), the systematic map protocol provided, as a first option, a real double coding for each article. In case of time constraints, it provided as a second option, that an a posteriori cross-check could be carried out with discussion between the coders on potential disagreements. Concerning the current project, due to the lack of means, we had to choose this alternative. We think that our approach did not affect coding consistency because (1) the coders followed the same rules, (2) a test of consistency was carried out before starting, with discussions on the potential disagreements, (3) all the reviewers were in contact throughout the coding

process, and (4) an a posteriori check was undertaken at the end of the coding step (see "Data coding strategy" section for more details).

Conclusions

This systematic map provides an up-to-date global catalogue of the available evidence on wildlife translocations to, from, or within protected areas. From the translocation synthesis, we identified a total of 498 publications equating to 841 translocations. Documented translocations were a result of various motivations but mainly driven by an aim to improve the conservation status of the focal species (it should also be noted that a large number of translocations did not mention any information on motivation). Translocation operations spanned a range of north and southern hemisphere regions (69 countries worldwide). Most studies aimed




Figure. 12. The number of translocations and their “direction” of transfer i.e. protected area context by taxonomic kingdom. “To”: transfer from a non-protected area (PA) to a PA; “From-To”: transfer from a PA to another; “Within same”: transfer occurring within the same PA perimeter; “From”: transfer from a PA to outside of PA. N.B., only known direction of movement is presented in the current figure





Fig. 13 Number of translocations by distance (in kilometres). N.B., This is only based on the number of translocations (358) where sufficient information was available in the publications to collate geographic coordinates either directly by authors or with detailed enough information given on capture and release sites to be able to retrieve coordinates manually

at assessing the survival, space use, demography, and reproduction of translocated species. To a lesser extent, the impacts on feeding habitats, behaviour, genetics, and physiology were also studied. The most translocated animal groups included mammals, birds, and reptiles, while the most translocated plant groups were *Magnoliopsida* and *Liliopsida*. The majority of species

are translocated to protected areas from non-protected sites.

Implications for policy/management

Given the map focused on translocations related to protected areas, the results are addressed primarily to their managers. The map database provides an estimate of the



extent of conservation translocations carried out worldwide, in which protected areas are involved, concerning all biodiversity (fauna, flora, fungi). In this respect, it should enable protected area managers to better perceive their role, in the global network of protected areas, both as suppliers or recipients of translocated organisms. Also, it could aid managers and practitioners in making their own choices by comparing previous experiences, regarding both the species concerned and the precise translocation modalities (number of individuals, etc.) (cf. Fig. 15 for a ‘walkthrough’ on how to navigate the database). Thus, at a time where reserve managers may be increasingly questioning whether to take such ‘risk-based’ actions, a detailed view of such interventions is clearly needed. As a consequence, the systematic map database provides valuable information to support decision making for future translocations. It may also be of interest to institutions and NGOs alike (i.e. Ministries, IUCN, etc.). If certain questions emerge from managers and decision makers, the map could also be used to develop one or more systematic reviews (for example to quantify the success factors involved in the survival of translocated individuals), which would further help guide decisions on future translocations.

Implications for research

The results of the current systematic map clearly identify that the evidence of translocations is skewed towards assessing survival (i.e. proportion of individuals alive or

level of mortality since translocation), demography (i.e. population growth overtime), and space use (i.e. the dispersal and home range distances). However, the impacts on the behavior, genetics, and physiology of translocated individuals will undoubtedly contribute to the long-term success of such interventions. Thus, we encourage researchers to study translocations using these outcomes to diversify the results. As a second point, this map identifies a number of groups that were associated with a few number of publications, such as *Anthozoa*, *Gastropoda*, *Bivalvia*, *Athropoda*, *Hexanauplia*, *Malacosraca*. This could be a lack of publications on existing translocations or a real absence of translocations for these species in question. It might be interesting to carry out further work to check whether certain translocations exist of which this map would therefore be unaware because they have not been published (which is in line with the limitations mentioned above on language restrictions, for example).

Lastly, concerning intervention type, 264/841 translocations were coded as *Unknown*. Indeed, the authors were unable to code with sufficient confidence whether the concerned translocations consisted of a true reintroduction, supplementation, or introduction. This was mainly due to either a lack of consistency or miss use of translocation vocabulary, or missing confirming information on in situ resident populations, or historical species’ distributions. We therefore underline the importance of using correct vocabulary and encourage future research to systematically publish details in the “Materials and

Flow diagram for managers navigating the evidence base – Additional File 7

Walkthrough: if I am a manager of an area considering translocating a population of mammals, and I require information from the scientific literature.




Fig. 15 illustrates how to navigate through the evidence base. Three walkthroughs are provided based on searching for information on mammal translocations. *N.B.*, these example questions are adapted from real operational inquiries addressed to the review team during workshops

Table 8 Comparing other evidence syntheses to our current map. N.B

Citation	Scope of review	Nature of synthesis	Search databases	No. of other literature sources	Publication data range of included articles	No. of included publications
<i>Our map (translocation synthesis)</i>	<i>P: all biodiversity worldwide I: reintroductions, introductions, Supplementations C: protected areas</i>	<i>Systematic map</i>	<i>2 databases: WOS, SCOPUS</i>	<i>12 (+2 grey literature calls)</i>	<i>1969 to 2020</i>	<i>498</i>
Fischer J, Lindenmayer DB. An assessment of the published results of animal relocations. <i>Biological Conservation</i> . 2000; 96: 1–11	P: animals worldwide I: reintroductions, supplementations, introductions	Map-like	0 databases	A search for articles across 12 major journals only	1979 to 1998	124
Hale SL, Koprowski JL. Ecosystem-level effects of keystone species reintroduction: a literature review. <i>Restoration Ecology</i> . 2018; 26: 439–45	P: key-stone species I: reintroductions only	Map-like	1 database: WOS	0	1995 to 2016	69
Tetzlaff SJ, Sperry JH, DeGregorio BA. Effects of antipredator training, environmental enrichment, and soft release on wildlife translocations: a review and meta-analysis. <i>Biol Cons</i> . 2019; 236: 324–31	P: all biodiversity ^a I: translocations ^a C: antipredator training, soft release, or environmental Enrichment	Meta-analysis	0 databases (Search in google scholar only)	0	1981 to 2018	41
Resende, P., Viana-Junior, A., Young, R., Azevedo, C., 2020. A global review of animal translocation programs. <i>Anim. Biodivers. Conserv.</i> 221–232. https://doi.org/10.32800/abc.2020.43.0221	P: animals I: introduction, reintroduction, translocations ^a	Map-like	2 databases: WOS, SCOPUS	1	1986 to 2017	145

P population, I interventions, C context

^a Methods unclear and exclusion criteria difficult to ascertain. The first line in italics corresponds to this map

methods” section of articles the exact context of a translocation intervention so that future map and review projects can incorporate as much metadata as possible.

Supplementary Information

The online version contains supplementary material available at <https://doi.org/10.1186/s13750-021-00236-w>.

Additional file 1. ROSES form.

Additional file 2. Search strategy.

Additional file 3. Test list and comprehensiveness.

Additional file 4. List of unobtained full text.

Additional file 5. Inclusion and exclusion decisions.

Additional file 6. List of accepted articles (all, unclear contexts, reviews/metapopulation management, translocation synthesis).

Additional file 7. List of studies (all, unclear contexts, link data, translocation synthesis).

Additional file 8. Calculating distances in R.

Additional file 9. List of rejected articles on language.

Acknowledgements

The authors would like to thank Dr. Hugo Mell for his contribution to full-text screening and coding; and Dr. Benoît Pisanu of the Natural heritage service (UMS Patrimoine Naturel OFB-CENRS-MNHN), notably for his expertise and knowledge on managing and analysing geographic data.

Authors' contributions

This systematic map is based on a protocol, Langridge et al. [60], written by JL, RS, and YR. JL and RS undertook searches, screening, and coding of articles. HM coded articles. JL and RS wrote the manuscript. All authors read and approved the final manuscript.

Authors' information

JL, RS, HM, YR are scientists.

Funding

This work was undertaken within the framework of the LIFE program entitled “NaturAdapt”. The project is co-financed by the LIFE programme (European commission), The French Ministry of Ecology and the French Office of Biodiversity.

Availability of data and materials

All data, generated or analysed during this systematic map are included in this published article and its Additional files.

Declarations

Ethics approval and consent to participate

Not applicable.

Consent for publication

Not applicable.

Competing interests

The authors declare having no competing interests.

Received: 15 March 2021 Accepted: 27 August 2021

Published online: 30 October 2021

References

- Vitousek PM, Mooney HA, Lubchenco J, Melillo JM. Human domination of earth's ecosystems. *Science*. 1997;277:494–9.
- Murphy SE, Greenaway F, Hill DA. Patterns of habitat use by female brown long-eared bats presage negative impacts of woodland conservation management. *J Zool*. 2012;288:177–83.
- Ceballos G, Ehrlich PR, Barnosky AD, García A, Pringle RM, Palmer TM. Accelerated modern human-induced species losses: entering the sixth mass extinction. *Sci Adv*. 2015;1:e1400253.
- Newbold T, Hudson LN, Hill SLL, Contu S, Lysenko I, Senior RA, et al. Global effects of land use on local terrestrial biodiversity. *Nature*. 2015;520:45–50.
- Nowakowski AJ, Thompson ME, Donnelly MA, Todd BD. Amphibian sensitivity to habitat modification is associated with population trends and species traits. *Glob Ecol Biogeogr*. 2017;26:700–12.
- IUCN. The IUCN global species programme. IUCN. 2015. <https://www.iucn.org/theme/species/about/iucn-global-species-programme>. Accessed 12 May 2021.
- Griffith B, Scott J, Carpenter J, Reed C. Translocation as a species conservation tool: status and strategy. *Science*. 1989;245:477–80.
- Chapin FS, Zavaleta ES, Eviner VT, Naylor RL, Vitousek PM, Reynolds HL, et al. Consequences of changing biodiversity. *Nature*. 2000;405:234–42.
- Rockström J, Steffen W, Noone K, Persson Å, Chapin FS, Lambin EF, et al. A safe operating space for humanity. *Nature*. 2009;461:472–5.
- Zari MP. Ecosystem services analysis in response to biodiversity loss caused by the built environment. SAPIENS Surveys and Perspectives Integrating Environment and Society. Institut Veolia Environnement; 2014. <http://journals.openedition.org/sapiens/1684>. Accessed 15 Feb 2021.
- Steffen W, Richardson K, Rockström J, Cornell SE, Fetzer I, Bennett EM, et al. Planetary boundaries: guiding human development on a changing planet. *Science*. 2015;347. <https://science.sciencemag.org/content/347/6223/1259855>. Accessed 12 May 2021.
- Dudley N, Shadie P, Stolton S. Guidelines for applying protected area management categories including IUCN WCPA best practice guidance on recognising protected areas and assigning management categories and governance types. IUCN; 2013. <https://portals.iucn.org/library/node/30018>. Accessed 18 May 2021.
- Cantú-Salazar L, Gaston K. Very large protected areas and their contribution to terrestrial biological conservation. *Bioscience*. 2010;60:808–18.
- Boucher TM, Spalding M, Revenga C. Role and trends of protected areas in conservation. In: Levin SA, editor. *Encyclopedia of biodiversity*, 2nd edition. Waltham: Academic Press; 2013. p. 485–503. <https://www.sciencedirect.com/science/article/pii/B9780123847195003488>. Accessed 12 May 2021.
- IUCN. Guidelines for reintroductions and other conservation translocations. IUCN; 2013. <https://portals.iucn.org/library/node/10386>. Accessed 12 May 2021.
- Turner IM, Corlett RT. The conservation value of small, isolated fragments of lowland tropical rain forest. *Trends Ecol Evol*. 1996;11:330–3.
- Schwartz MW, van Mantgem PJ. The value of small preserves in chronically fragmented landscapes. In: Schwartz MW, editor. *Conservation in highly fragmented landscapes*. Boston: Springer; 1997. p. 379–94. https://doi.org/10.1007/978-1-4757-0656-7_16.
- IUCN. Guidelines for reintroductions and other conservation translocations. IUCN; 2013. <https://www.iucn.org/content/guidelines-reintroductions-and-other-conservation-translocations>. Accessed 31 Oct 2019.
- Seddon PJ, Griffiths CJ, Soorae PS, Armstrong DP. Reversing defaunation: restoring species in a changing world. *Science*. 2014;345:406–12.
- Cheyne S. Wildlife reintroduction: considerations of habitat quality at the release site. *BMC Ecol*. 2006;6:5.
- Moorhouse TP, Gelling M, Macdonald DW. Effects of habitat quality upon reintroduction success in water voles: evidence from a replicated experiment. *Biol Conserv*. 2009;142:53–60.
- Godefroid S, Piazza C, Rossi G, Buord S, Stevens A-D, Aguruija R, et al. How successful are plant species reintroductions? *Biol Conserv*. 2011;144:672–82.
- Stadtman S, Seddon PJ. Release site selection: reintroductions and the habitat concept. *Oryx*. 2020;54:687–95.

24. Peters RL, Darling JDS. The greenhouse effect and nature reserves global warming would diminish biological diversity by causing extinctions among reserve species. *Bioscience*. 1985;35:707–17.
25. Seddon PJ. From reintroduction to assisted colonization: moving along the conservation translocation spectrum. *Restor Ecol*. 2010;18:796–802.
26. Kostyack J, Lawler JJ, Goble DD, Olden JD, Scott JM. Beyond reserves and corridors: policy solutions to facilitate the movement of plants and animals in a changing climate. *Bioscience*. 2011;61:713–9.
27. Lunt ID, Byrne M, Hellmann JJ, Mitchell NJ, Garnett ST, Hayward MW, et al. Using assisted colonisation to conserve biodiversity and restore ecosystem function under climate change. *Biol Conserv*. 2013;157:172–7.
28. Watson J, Dudley N, Segan D, Hockings M. The performance and potential of protected areas. *Nature*. 2014;515:67–73.
29. Heller NE, Zavaleta ES. Biodiversity management in the face of climate change: a review of 22 years of recommendations. *Biol Conserv*. 2009;142:14–32.
30. Prober SM, Doerr VAJ, Broadhurst LM, Williams KJ, Dickson F. Shifting the conservation paradigm: a synthesis of options for renovating nature under climate change. *Ecol Monogr*. 2019;89:e01333.
31. Sandom CJ, Dempsey B, Bullock D, Ely A, Jepson P, Jimenez-Wisler S, et al. Rewilding in the English uplands: policy and practice. *J Appl Ecol*. 2019;56:266–73.
32. Macias-Fauria M, Jepson P, Zimov N, Malhi Y. Pleistocene Arctic megafaunal ecological engineering as a natural climate solution? *Philos Trans R Soc B Biol Sci*. 2020;375:20190122.
33. Seddon PJ, Armstrong DP, Maloney RF. Developing the science of reintroduction biology. *Conserv Biol*. 2007;21:303–12.
34. Seddon P, Armstrong D. 2. Reintroduction and other conservation translocations: history and future developments. In: *Reintroduction of fish and wildlife populations*. Oakland: University of California Press; 2019. p. 7–28.
35. Abeli T, Cauzzi P, Rossi G, Pistoja F, Mucciarelli M. A gleam of hope for the critically endangered *Isoetes malinverniana*: use of small-scale translocations to guide conservation planning. *Aquat Conserv Mar Freshw Ecosyst*. 2018;28:501–5.
36. Jolly CJ, Kelly E, Gillespie GR, Phillips B, Webb JK. Out of the frying pan: reintroduction of toad-smart northern quolls to southern Kakadu National Park. *Austral Ecol*. 2018;43:139–49.
37. Briers-Louw WD, Verschuere S, Leslie AJ. Big cats return to Majete wildlife reserve, Malawi: evaluating reintroduction success. *Afr J Wildl Res*. 2019;49:34–50.
38. Hitchcock A, Williams J, Cowell C. Lessons learned as *Erica turgida* is returned. *J Nat Conserv*. 2020;56:125858.
39. Moehrenschrager A, Macdonald DW. Movement and survival parameters of translocated and resident swift foxes *Vulpes velox*. *Anim Conserv Forum*. 2003;6:199–206.
40. Clements D. Translocation of rare plant species to restore Garry oak ecosystems in western Canada: challenges and opportunities. *Botany*. 2013;91:283–91.
41. Jackson CL, Schuster R, Arcese P. Release date influences first-year site fidelity and survival in captive-bred Vancouver Island marmots. *Ecosphere*. 2016;7:e01314.
42. Steenweg R, Hebblewhite M, Gummer D, Low B, Hunt B. Assessing potential habitat and carrying capacity for reintroduction of plains bison (*Bison bison*) in Banff National Park. *PLoS ONE*. 2016;11:e0150065.
43. Sarrazin F, Barbault R. Reintroduction: challenges and lessons for basic ecology. *Trends Ecol Evol*. 1996;11:474–8.
44. McCoy ED, Osman N, Hauch B, Emerick A, Mushinsky HR. Increasing the chance of successful translocation of a threatened lizard. *Anim Conserv*. 2014;17:56–64.
45. Fischer J, Lindenmayer DB. An assessment of the published results of animal relocations. *Biol Conserv*. 2000;96:1–11.
46. Azar JF, Rautureau P, Lawrence M, Calabuig G, Hingrat Y. Survival of reintroduced Asian houbara in United Arab Emirates' reserves. *J Wildl Manag*. 2016;80:1031–9.
47. Bricchieri-Colombi TA, Moehrenschrager A. Alignment of threat, effort, and perceived success in North American conservation translocations. *Conserv Biol*. 2016;30:1159–72.
48. Morris SD, Brook BW, Moseby KE, Johnson CN. Factors affecting success of conservation translocations of terrestrial vertebrates: a global systematic review. *Glob Ecol Conserv*. 2021;28:e01630.
49. Heiken KH, Brusck GA, Gartland S, Escallón C, Moore IT, Taylor EN. Effects of long distance translocation on corticosterone and testosterone levels in male rattlesnakes. *Gen Comp Endocrinol*. 2016;237:27–33.
50. de Milliano J, Stefano JD, Courtney P, Temple-Smith P, Coulson G. Soft-release versus hard-release for reintroduction of an endangered species: an experimental comparison using eastern barred bandicoots (*Perameles gunnii*). *Wildlife*. 2016;43:1–12.
51. Tetzlaff S, Sperry J, Degregorio B. Effects of antipredator training, environmental enrichment, and soft release on wildlife translocations: a review and meta-analysis. *Biol Conserv*. 2019;236:324–31.
52. Corlett RT. Restoration, reintroduction, and rewilding in a changing world. *Trends Ecol Evol*. 2016;31:453–62.
53. Mukesh, Sharma LK, Charoo SA, Sathyakumar S. Conflict bear translocation: investigating population genetics and fate of bear translocation in Dachigam National Park, Jammu and Kashmir, India. *PLoS ONE*. 2015;10:e0132005.
54. Harvey D, Lentini A, Cedar K, Weatherhead P. Moving massasaugas: insight into rattlesnake relocation using *Sistrurus c. catenatus*. *Herpetol Conserv Biol*. 2014;9:67–75.
55. Hale SL, Koprowski JL. Ecosystem-level effects of keystone species reintroduction: a literature review. *Restor Ecol*. 2018;26:439–45.
56. Resende P, Viana-Junior A, Young R, Azevedo C. A global review of animal translocation programs. *Anim Biodivers Conserv*. 2020;43:221–32.
57. McLachlan JS, Hellmann JJ, Schwartz MW. A framework for debate of assisted migration in an era of climate change. *Conserv Biol*. 2007;21:297–302.
58. Weeks AR, Sgro CM, Young AG, Frankham R, Mitchell NJ, Miller KA, et al. Assessing the benefits and risks of translocations in changing environments: a genetic perspective. *Evol Appl*. 2011;4:709–25.
59. Letty J, et al. Improving translocation success: an experimental study of anti-stress treatment and release method for wild rabbits. *Anim Conserv Forum*. 2000;3:211–9.
60. Langridge J, Sordello R, Reyjol Y. Outcomes of wildlife translocations in protected areas: what is the type and extent of existing evidence? A systematic map protocol. *Environ Evid*. 2020;9:16. <https://doi.org/10.1186/s13750-020-00199-4>.
61. Guidelines for authors—environmental evidence. <https://environmentalevidence.org/information-for-authors/>. Accessed 22 Feb 2021.
62. Haddaway NR, Collins AM, Coughlin D, Kirk S. The role of google scholar in evidence reviews and its applicability to grey literature searching. *PLoS ONE*. 2015;10:e0138237. <https://doi.org/10.1371/journal.pone.0138237>.
63. James KL, Randall NP, Haddaway NR. A methodology for systematic mapping in environmental sciences. *Environ Evid*. 2016;5:7.
64. Fernandez FAS, Rheingantz ML, Genes L, Kenup CF, Galliez M, Cezimbra T, et al. Rewilding the Atlantic Forest: restoring the fauna and ecological interactions of a protected area. *Perspect Ecol Conserv*. 2017;15:308–14.
65. Bowler DE, Bjorkman AD, Dornelas M, Myers-Smith IH, Navarro LM, Niamir A, et al. Mapping human pressures on biodiversity across the planet uncovers anthropogenic threat complexes. *People Nat*. 2020;2:380–94.
66. Zimmer H, Auld T, Cuneo P, Offord C, Commander L. Conservation translocation—an increasingly viable option for managing threatened plant species. *Aust J Bot*. 2019;67:501–9.
67. Kottek M, Grieser J, Beck C, Rudolf B, Rubel F. World Map of the Köppen-Geiger climate classification updated. *Meteorol Z*. 2006;15:259–63.
68. Beck HE, Zimmermann NE, McVicar TR, Vergopolan N, Berg A, Wood EF. Present and future Köppen-Geiger climate classification maps at 1-km resolution. *Sci Data*. 2018;5:180214.
69. Zarrouk A, Romdhane MS, Espinosa F. Reintroduction of the highly endangered mollusk *Patella ferruginea* Gmelin, 1791 in an MPA: a novel approach to achieve high survival rates. *Estuar Coast Shelf Sci*. 2018;202:259–69.
70. Germano J, Bishop P. Suitability of amphibians and reptiles for translocation. *Conserv Biol*. 2009;23:7–15.
71. Tosi G, Chirichella R, Zibordi F, Mustoni A, Giovannini R, Groff C, et al. Brown bear reintroduction in the Southern Alps: to what extent are expectations being met? *J Nat Conserv*. 2015;26:9–19.
72. Bennett VA, Doerr VAJ, Doerr ED, Manning AD, Lindenmayer DB, Yoon H-J. Habitat selection and post-release movement of reintroduced brown treecreeper individuals in restored temperate woodland. *PLoS ONE*. 2012;7:e50612.

73. Muller LI, Murrow JL, Lupardus JL, Clark JD, Yarkovich JG, Stiver WH, et al. Genetic structure in Elk persists after translocation. *J Wildl Manag*. 2018;82:1124–34.
74. Watts C, Thornburrow D. Where have all the weta gone? Results after two decades of transferring a threatened New Zealand giant weta, *Deinacrida mahoenui*. *J Insect Conserv*. 2008;13:287–95.
75. Goldenberg SZ, Owen MA, Brown JL, Wittemyer G, Oo ZM, Leimgruber P. Increasing conservation translocation success by building social functionality in released populations. *Glob Ecol Conserv*. 2019;18:e00604.
76. Tarszisz E, Dickman CR, Munn AJ. Physiology in conservation translocations. *Conserv Physiol*. 2014;2. <https://www.ncbi.nlm.nih.gov/pmc/articles/PMC4732500/>. Accessed 14 June 2021.
77. Kenup CF, Sepulveda R, Kreischer C, Fernandez FAS. Walking on their own legs: unassisted population growth of the agouti *Dasyprocta leporina*, reintroduced to restore seed dispersal in an Atlantic Forest reserve. *Oryx*. 2018;52:571–8.
78. Bennett VA, Doerr VAJ, Doerr ED, Manning AD, Lindenmayer DB. The anatomy of a failed reintroduction: a case study with the Brown Treecreeper. *Emu Austral Ornithol*. 2012;112:298–312.
79. Baker J, Bain D, Clarke J, French K. Translocation of the Eastern Bristlebird 2: applying principles to two case studies. *Ecol Manag Restor*. 2012. <https://doi.org/10.1111/j.1442-8903.2012.00640.x>.
80. Muths E, Johnson T, Corn P. Experimental repatriation of boreal toad (*Bufo boreas*) eggs, metamorphs, and adults in Rocky Mountain National Park. *Southwest Nat*. 2001;46:106.

Publisher's Note

Springer Nature remains neutral with regard to jurisdictional claims in published maps and institutional affiliations.

Ready to submit your research? Choose BMC and benefit from:

- fast, convenient online submission
- thorough peer review by experienced researchers in your field
- rapid publication on acceptance
- support for research data, including large and complex data types
- gold Open Access which fosters wider collaboration and increased citations
- maximum visibility for your research: over 100M website views per year

At BMC, research is always in progress.

Learn more biomedcentral.com/submissions

